An Update on CLEO's Study of B Meson Decays

Bernard Gittelman

Cornell University - Ithaca New York, USA

Introduction

The Υ family of mesons was discovered at Fermilab[1] in 1977. At that time, purely by coincidence the CESR storage ring had been designed and was under construction. Construction of the CLEO-I detector had also begun. In 1978, the existence of the $\Upsilon(1S)$ and $\Upsilon(2S)$ resonances were confirmed at the DORIS storage ring of DESY[2], and from their measurement of the $e^+e^- \to \Upsilon(1S)$ crossection, it became clear that the Υ resonances are best described as $b\bar{b}$ quark pairs, not $t\bar{t}$. CESR began operation at the end of 1979. The CLEO (and CUSB) Collaborations began data taking. In 1980, CLEO (and CUSB) published observation of the $\Upsilon(1S)$, $\Upsilon(2S)$, $\Upsilon(3S)$[3], and the $\Upsilon(4S)$[4]. The $\Upsilon(4S)$, is a much wider resonance and is believed to decay exclusively to $B\bar{B}$ mesons. Over the next six to eight years, many general properties of B meson decay were measured at DORIS and CESR. In 1988, CESR shut down and the CLEO-I detector was replaced by the CLEO-II detector. The new CLEO-II detector consists of a better resolution tracking system and electromagnetic calorimeter. Figure 1 is an illustration of the CLEO-II detector. Our understanding of the B meson decay has continued to improve with the CLEO-II detector, but today we only understand approximately 50% of the B decays. This paper is a description of the status of our understanding today compared with the Particle Data Group review published in 1996.

Understanding of B Meson Decay in 1996

In the 1996 Review of Particle Physics[5], the Particle Data Group pro-

vided a long detailed summary of what is known about the B meson branching fractions, and life times. Most of the information they reported on the B meson branching fractions came from publications of the ARGUS Collaboration at DESY and publications of CLEO at Cornell. A summary of the "largest" B meson branching fractions listed in the 1996 Review of Particle Physics is given in Table 1. In the Standard Model, the b quark decays under the weak interaction to the c or u quark. From early experiments, it was evident that the $b \to u$ interaction is more than an order of magnitude weaker than the $b \to c$ (See for example the $b \to u l \nu$ and the $B \to X_c l \nu$ branching fractions listed in Table 1.). B meson decays are usually described by the diagrams shown in figure 2 (Since the main decays are to c quarks, the diagrams shown in figure 2 do not include the $b \to u$ transitions.). The first set of branching fractions in Table 1 are the semileptonic decay modes. These are the simplest decay modes to describe. They all proceed via the tree diagram. The listed branching fractions, for example $Br(B \to X l \nu) = (10.4 \pm 0.4)\%$, is supposed to represent the branching fraction for $l = e$ or $l = \mu$. The total semileptonic branching fraction includes $B \to X e \nu, B \to X \mu \nu$, and $B \to X \tau \nu$. The branching fraction to electons and muons are expected to be very close to identical. However, the branching fraction to τ leptons, which have never been measured, are expected to be only 0.3 as large, because of the large τ mass (The factor 0.3 comes from the phase space). The sum over electron, muon and tau semileptonic decay is $2.3 \times (10.4 \pm 0.4) = (23.9 \pm 0.9)\%$.

The second group of branching fractions in Table 1 are the 2 body decays of the B meson in which the W boson materializes into a $u\bar{d}$, (i.e., π^+ or ρ^+). One notices that the B^+ branching fractions are 1.5 - 2 times larger than the B^o's. The diagrams describing the $B^+ \to \pi^+ D^o$ decay are the tree and the color supressed shown in figure 2. The $B^o \to \pi^+ D^-$ decay proceeds through the tree and the W exchange diagram. If the W exchange amplitude is as large as the color supressed amplitude, this would imply that color supressed amplitude is in phase with the tree, and the W exchange amplitude is out of phase.

The third group of branching fractions in Table 1 are the 2 body decays of the B meson into a $D^{(*)} D_s^{(*)}$. These exclusive two body modes add up to $\approx (5.5 \pm 2.0)\%$. The inclusive yield of D_s coming from B decays was listed as $(8.6 \pm 2.2)\%$, which implies 64% of the $B \to X D_s$ are two body.

The inclusive decay of B mesons to $\bar{D}$ mesons in Table 1 add up to $(82.2 \pm 6.0)\%$. These include the D mesons coming from semileptonic decay

of the B mesons. Most of the rest of B meson decays are into Charmonium or Charm-Baryons. Adding the 1.14 ± 0.06 and the 6.4 ± 1.1 to the $82.2 \pm 6.0\%$, one has accounted for $(89.7 \pm 6.1)\%$ of B meson decays.

Inclusive B Meson Decay Branching Fractions

During the past few years, CLEO has made many remeasurements of inclusive branching fractions in B meson decay. The most recent CLEO measurements on the inclusive branching fractions are listed in Table 2. These results are generally consistent but a little more accurate than those shown in Table 1. The separate measurement of semileptonic decays of B^+ and B^o mesons was first made via full reconstruction of the $\bar{B}$ meson hadronic decay in the event then examining how often there was a lepton coming from the remaining B meson decay[6]. Although this method was very clear and simple, it was statistically limited by the small sample of fully reconstructed B (or $\bar{B}$) mesons. The results listed in Table 2 were obtained by partial reconstruction of $\bar{B}^o \to D^{*+} l^- \nu, D^{*+} \to \pi^+ D^o$, detecting only the lepton and the π^+[7]. Having collected a very large sample of partially reconstructed B^os and measuring how often there is also a lepton amongst the leftover tracks coming from the other B meson decay, the semileptonic branching fraction of the B^o was determined. To obtain the $B^- \to X^o l^- \nu$ branching fraction, the result of this analysis was then combined with a previous analysis[13] in which the semileptonic branching fraction averaged over charged and neutral B mesons was measured. The result indicates the B^o branching fraction is 0.53% larger than the B^+, which is indicates the B^o lifetime is 0.53% longer than the B^+. However, the uncertainty is large enough so that the branching fractions could be equal.

The inclusive decays to D mesons in table 2 are the result of observing how often it was possible to reconstruct a D^o (via $D^o \to K^- \pi^+$) and how often it was possible to reconstruct D^+ (via $D^+ \to K^- \pi^+ \pi^+$) in a set of events coming from $\Upsilon(4S)$ decay. The same analysis was used to measure the inclusive B branching fraction to D^{*+} and D^{*o} by measuring how often it is possible to reconstruct a D^{*+} and how often a D^{*o}. One should understand the inclusive D branching fractions include D mesons coming from D^* decays.

Normally, the D mesons from $\bar{B}$ meson decay originate from the b quark transition, $b \to cW^-$. However, $\bar{D}$ mesons can come from $\bar{B}$ meson decay by the $W^- \to \bar{c}s$. Both the tree and the color supressed diagrams (see figure 2)

Table 1.
Summary of Major B Meson Branching Fractions known in 1996

(Numbers from PDG Review[5])

Decay Mode	B^+ (%)	B^o (%)	**Admixture** B^+ **and** B^o (%)
$B \to \bar{D} l\nu$	1.6 ± 0.7	1.9 ± 0.5	
$B \to \bar{D}^* l\nu$	5.3 ± 0.8	4.6 ± 0.3	
$B \to \bar{D}^{**} l\nu$			2.7 ± 0.7
$B \to X_c l\nu$	10.1 ± 2.3	10.3 ± 1.0	
$B \to X l\nu$			10.4 ± 0.4
$b \to u l\nu$			$\sim 0.18 \pm 0.05$
$B \to \bar{D} \rho^+$	1.34 ± 0.18	0.78 ± 0.14	
$B \to \bar{D}^* \rho^+$	1.55 ± 0.31	0.73 ± 0.15	
$B \to \bar{D} \pi^+$	0.53 ± 0.05	0.30 ± 0.04	
$B \to \bar{D}^* \pi^+$	0.52 ± 0.08	0.26 ± 0.04	
$B \to \bar{D} D_s^+$	1.7 ± 0.6	0.7 ± 0.4	
$B \to \bar{D}^* D_s^+$	0.1 ± 0.07	1.2 ± 0.6	
$B \to \bar{D} D_s^{*+}$	1.2 ± 1.0	2.0 ± 1.5	
$B \to \bar{D}^* D_s^{*+}$	2.3 ± 1.4	1.9 ± 1.2	
$B \to X D_s^+$			8.6 ± 2.2
$B \to D^- X$			24.2 ± 3.3
$B \to \bar{D}^o X$			58.0 ± 5.0
$B \to \Psi K$	0.10 ± 0.014	0.075 ± 0.021	
$B \to \Psi K^+ \pi^+ \pi^-$	0.14 ± 0.06		
$B \to \Psi K^+ \pi^-$		0.11 ± 0.06	
$B \to \Psi K^*$	0.17 ± 0.05	0.16 ± 0.03	
$B \to \Psi X$			1.14 ± 0.06
$B \to$ **Charm-Baryons**			6.4 ± 1.1

can lead to $\bar{D}$ from $\bar{B}$ decays. A measurement of how often the $\bar{B}$ meson decays to a D meson and how often to a $\bar{D}$ meson has been made[9]. This was carried out by using the fact that the $\Upsilon(4S)$ decays to $B\bar{B}$. One of the two B mesons is tagged via the sign of the lepton in its semileptonic decay. Only high momentum leptons ($p_l > 1.4$ GeV/c) are used, to eliminate lepton from semileptonic D decay. The other B meson is assumed to have the opposite flavor (Corrections are made for the $B^o \leftrightarrow \bar{B}^o$ mixing of the neutral B mesons). Having tagged the flavor of one of the B mesons, the D meson coming from that same B meson is known to have an explicit flavor. This D meson, which comes from the tagged B meson, is identified via the angular correlation. Since the lepton has a momentum greater than 1.4 GeV/c, the D meson must be moving at an angle greater than 90^o away from the lepton in order to conserve momentum. The D mesons from the other B meson are isotropic, so the flavor of the D mesons in the same hemisphere as the lepton was studied. From the numbers in Table 2, one sees that 90% of the D mesons from B decay come from the c quark that originates from the $b \to cW^-$ transition and 10% come from the W boson.

The $B \to XD_s$ inclusive branching fractions were measured by studying the number of D_s mesons and their momentum spectrum at the $\Upsilon(4S)$ No attempt was made to tag the flavor of the B meson or its charge. From the D_s momentum spectrum it was concluded that the D_s is almost always coming from the W boson. To obtain the branching fraction, 12.11%, it was assumed that the B^+ and B^o branching fraction to D_s are the same. This value is significantly larger than the 1996 PDG value (see Table 1.).

The most interesting new inclusive result is the measurement of the B decay to ηX[12]. The 17.6% branching fraction is much larger than one might expect. Some of the η mesons come from the D and D_s mesons that are in the B meson decay. The contribution from feed down has been estimated from the inclusive $B \to \bar{D}^o, D^-$, and D_s branching fractions and the inclusive η branching fractions of each of these three charm mesons. The branching fraction estimates are $\mathcal{B}r(B \to \bar{D}X \to \eta X') = (2.5 \pm 0.5)\%$, and $\mathcal{B}r(B \to D_s X \to \eta X') = (2.8 \pm 0.6)\%$. Subtracting these leads to the conclusion $(12.3 \pm 1.8)\%$ of B decays produce η mesons directly. η mesons are often described as partly an $s\bar{s}$ resonance, but this 12.3% branching fraction is much larger than the incusive $\mathcal{B}r(B \to \phi X) = (3.5 \pm 0.7)\%$[5].

Table 2,
Inclusive B Meson Branching Fractions from CLEO

Decay Mode	**Branching Fraction (%)**	**Reference**
Semileptonic Decays		
$B^+ \to X^o l^+ \nu$	$10.25 \pm 0.57 \pm 0.65$	[**7**]
$B^o \to X^- l^+ \nu$	$10.78 \pm 0.60 \pm 0.69$	[**7**]
Inclusive Decays to D mesons		
Assuming an Equal Admixture of $B^o\bar{B}^o$ and B^+B^-		
$B \to D^o X + \bar{B} \to D^o X$	$63.6 \pm 1.4 \pm 2.6$	[**8**]
$B \to D^+ X + \bar{B} \to D^+ X$	$23.5 \pm 0.9 \pm 2.4$	[**8**]
$B \to D^{*o} X + \bar{B} \to D^{*o} X$	$24.7 \pm 1.2 \pm 2.5$	[**8**]
$B \to D^{*+} X + \bar{B} \to D^{*+} X$	$23.9 \pm 1.5 \pm 1.7$	[**8**]
Inclusive Flavor Sensitive Decays to D mesons		
(lv) for D meson from $b \to c$, (uv) for D meson from $W^- \to \bar{c}s$		
$\bar{B} \to D^o X + \bar{B} \to D^+ X$ **(lv)**	79.2 ± 3.9	[**9**]
$\bar{B} \to \bar{D}^o X + \bar{B} \to D^- X$ **(uv)**	7.9 ± 2.2	[**9**]
Inclusive B $\to D_s$ Decays		
$B^+ \to D_s^+ X = B^o \to D_s^+ X$	$12.11 \pm 0.39 \pm 1.63$	[**10**]
Inclusive B $\to$ Charmonium Decays		
$B \to \Psi X = \bar{B} \to \Psi X$	$1.12 \pm 0.04 \pm 0.06$	[**11**]
$B \to \Psi X = \bar{B} \to \Psi X$ **Direct**	0.80 ± 0.08	[**11**]
$B \to \Psi' X = \bar{B} \to \Psi' X$	$0.34 \pm 0.04 \pm 0.03$	[**11**]
$B \to \chi_{c1} X = \bar{B} \to \chi_{c1} X$	$0.40 \pm 0.06 \pm 0.04$	[**11**]
$B \to \chi_{c2} X = \bar{B} \to \chi_{c2} X$	$0.25 \pm 0.10 \pm 0.03$	[**11**]
$B \to$ **Charmonium (total)**	1.79 ± 0.15	**from above**
Inclusive B Decays to ηX		
$(B \to \eta X)$	$17.6 \pm 1.1 \pm 1.2$	[**12**]

Exclusive B Meson Decay Branching Fractions

The sum of all the exclusive branching fractions listed in Table 1 add up to $(15.6 \pm 2.1)\%$, (16.6% for B^+, 17.4% for B^o). In recent years, many other 2 body final states have been searched for. Most of these are "rare B decays"[14]. The exclusive final states listed in Table 1 have been remeasured. The new CLEO results for these and a few others are given in Table 3. There have been several new measurements of semileptonic decays. For the $b \to c$ final states, there have been no serious changes. The $D_1^o l\nu$ has been measured. There have also been measurements of the $b \to u$ final states, $\pi l\nu$ and $\rho l\nu$. The branching fractions of the exclusive $b \to cl\nu$ final states that have been observed add up to 7.0% which is only $\sim$ 67% of the inclusive branching fraction. The observed $b \to u$ final states add up to 0.043% compared with the inclusive rate of 0.18% from Table 1, i.e., only $\sim$ 25% of the $b \to ul\nu$ is understood.

The exclusive two body charm decays in which the W boson materializes as a π, ρ or D_s have been remeasured. Most of the results agree with the earlier measurements and are more accurate. The remeasurements on $B \to \Psi K^{(*)}$ have similar results.

Summary- What Fraction of B Meson Decays are "Understood"

Theoretically, the Semileptonic decays of B mesons are the easiest to understand, because there is only one hadronic vertex. Originally[25], the semileptonic branching fraction was estimated to be approximately 16%. After independent measurements at DORIS, CESR, and LEP, we find the branching fraction is significantly smaller. The generic average of the CLEO measurements gives the inclusive branching fraction,

$$\mathcal{B}r(\bar{B} \to Xl\nu) = (10.5 \pm 0.89)\%.$$

Although using QCD corrections, theorists have recently suggested the B semileptonic branching fraction could be as low as 12%, the CLEO result is a theoretical challenge. Furthermore, only a fraction of the of the hadronic content of the semileptonic decays have been measured. Summing over the

Table 3,
Exclusive B meson Decay Branching Fractions from CLEO

Decay Mode	B^+ (%)	B^o (%)	**Reference**
Semileptonic Decays			
$B \to Dl\nu$	$1.94 \pm 0.15 \pm 0.34$	$1.87 \pm 0.15 \pm 0.32$	[**15**]
$B \to D^* l\nu$	$5.13 \pm 0.54 \pm 0.64$	$4.49 \pm 0.32 \pm 0.39$	[**16**]
$B \to D_1^o l\nu$	$0.56 \pm 0.13 \pm 0.09$	**?**	[**17**]
$B \to \pi l\nu$		$0.018 \pm 0.004 \pm 0.004$	[**18**]
$B \to \rho l\nu$		$0.025 \pm 0.004 \pm 0.007$	[**18**]
Exclusive Two Body Charm Decays			
$B \to D\rho$	$1.35 \pm 0.12 \pm 0.15$	$0.81 \pm 0.11 \pm 0.18$	[**19**]
$B \to D^*\rho$	$1.68 \pm 0.21 \pm 0.28$	$0.74 \pm 0.10 \pm 0.14$	[**19**]
$B \to D\pi$	$0.55 \pm 0.04 \pm 0.05$	$0.29 \pm 0.04 \pm 0.06$	[**19**]
$B \to D^*\pi$	$0.434 \pm 0.033 \pm 0.038$	$0.281 \pm 0.011 \pm 0.022$	[**20**]
$B \to DD_s^+$	$1.26 \pm 0.22 \pm 0.29$	$0.87 \pm 0.24 \pm 0.22$	[**21**]
$B \to DD_s^{*+}$	$0.87 \pm 0.27 \pm 0.20$	$1.00 \pm 0.35 \pm 0.25$	[**21**]
$B \to D^* D_s^+$	$1.40 \pm 0.43 \pm 0.39$	$0.93 \pm 0.23 \pm 0.19$	[**21**]
$B \to D^* D_s^{*+}$	$3.10 \pm 0.88 \pm 0.75$	$2.03 \pm 0.50 \pm 0.43$	[**21**]
$B \to D^- D^+$		< 0.12	[**22**]
$B \to D^{*\pm} D^{\mp}$		< 0.18	[**22**]
$B \to D^{*-} D^{*+}$		< 0.22	[**22,26**]
Exclusive B $\to$ Charmonium Decays			
$B \to \Psi K$	$0.102 \pm 0.008 \pm 0.007$	$0.085 \pm 0.013 \pm 0.006$	[**23**]
$B \to \Psi K^*$	$0.141 \pm 0.023 \pm 0.024$	$0.132 \pm 0.017 \pm 0.017$	[**23**]
$B \to \Psi\pi$	0.0056 ± 0.0027	< 0.0058	[**24**]
$B \to \Psi\rho$	< 0.077	< 0.025	[**24**]
$B \to \Psi\omega$		< 0.027	[**24**]

measured charm hadronic states, one has

$$\mathcal{B}r(\bar{B} \to (D + D^* + D^{**})l\nu) = (7.00 \pm 0.85)\%$$

Therefore, we understand $(7.00 \pm 0.85)/(10.5 \pm 0.89) = (67.0 \pm 13.7)\%$ of semileptonic branching fractions.

From the CLEO measurement of the flavor dependence of the decay of B mesons to D mesons, we know the $\mathcal{B}r(\bar{B} \to DX) = (79.2 \pm 3.9)\%$ (i.e., from $b \to c$ at the lower vertex). Assuming the exclusive semileptonic branching fractions to τ's are 0.3 times the branching fractions to electrons and/or muons, and adding the exclusive semileptonic decays to e's, μ's, and τ's, we have for the total exclusive semileptonic branching fraction

$$\mathcal{B}r(\bar{B} \to D^{(*,**)}(e + \mu + \tau)\nu) = (2.3 \times (7.00 \pm 0.85) = (16.1 \pm 2.0)\%$$

The exclusive two body decays resulting in a D meson add up to

$$\mathcal{B}r(\bar{B} \to D\pi + D^*\pi + D\rho + D^*\rho + DD_s + D^*D_s + DD_s^* + D^*D_s^*) = (8.8 \pm 1.2)\%$$

(This was calculated from the numbers in Table 3, averaging over B^+ and B^o.) Putting the above numbers together, the inclusive $B \to DX$ decays are "understood" at the level,

$$\mathcal{B}r(\bar{B} \to D^{(*)}(l\nu + \pi + \rho + D_s)) = (16.1 \pm 2.0) + (5.73 \pm 1.17) + (3.07 \pm 0.35)$$
$$= (24.9 \pm 1.8)\%$$

This is $(24.9 \pm 1.8)/(79.2 \pm 3.9) = 0.31 \pm 0.03$ of the inclusive D mesons coming from the $b \to c$ vertex.

The average of exclusive final states with charmonium that have been measured add up to $\sim (0.23 \pm 0.03)\%$. The exclusive baryon final states have not yet been fully reconstructed. Therefore, we have an exclusive understanding of $(24.9 \pm 1.8 + 0.23 \pm 0.03) = (25.1 \pm 1.8)\%$ of the B decays.

In terms of inclusive branching fractions, in the $\bar{B} \to DX$ decays, the X represents $l\nu X'$, $D_s X''$, and X'''. Here X' includes the π's coming from D^* or D^{**} which we know something about, but X'' and X''' are not known. From the value of the branching fraction $\bar{B} \to DX = (79.2 \pm 3.9)\%$, we calculate,

$$\mathcal{B}r(\bar{B} \to DX''') = \bar{B} \to D(X - l\nu X' - D_s X'')$$
$$= (79.2 \pm 3.9) - (2.3 \times (10.5 \pm 0.89)) - (12.1 \pm 1.7)$$
$$= (42.9 \pm 4.7)\%$$

This means we know nothing about $(42.9/79.2) = 54\%$ of the inclusive $B \to DX$ decays.

The total inclusive branching fractions, (excluding the "rare" decays) add up to

$$\begin{aligned} \mathcal{B}r(B \to DX + charmoniumX' + baryonsX'') = \\ = \quad (79.2 \pm 3.9) + (1.8 \pm 0.2) + (6.4 \pm 1.1) \\ = \quad (87.4 \pm 4.1)\% \end{aligned}$$

CLEO has recently made many measurements of rare decays (and upper limits of rare decays)[14]. However, the total sum of the measured modes and the upper limits on rare B decay branching fractions is less than 2×10^{-3}, so there is no need to includes these at this time. The conclusion is one must invent non-rare decay modes and look for them.

Acknowledgements

I would like to thank Giancarlo Moneti and Ed Thorndike of the CLEO Collaboartion for useful comments, corrections, and discussions related to this paper. The hospitality of the organizers of the Quarks-98 Seminar in Suzdal made this visit to Russia, my first in ten years, warm and interesting.

References

[1] Observation of a new group of resonances at 10 GeV decaying to muons pairs
S.W. Herb et al., Phys. Rev. Lett. 39, 252(1977).

[2] Confirmation of the existence of the $\Upsilon(1S)$ and $\Upsilon(2S)$
C. Berger et al., Phys. Lett. 76*B*, 243(1978).
C.W. Darden et al., Phys. Lett. 78*B*, 246(1978).
J.K. Bienlein et al., Phys. Lett. 78*B*, 360(1978).
C.W. Darden et al., Phys. Lett. 80*B*, 419(1979).

[3] Observation of Three Upsilon States
D. Andrews et al., Phys. Rev. Lett. 44, 1108(1980).
T. Bohringer et al., Phys. Rev. Lett. 44, 1111(1980).

[4] Observation of a Fourth Upsilon State in e^+e^- Annihilations
D. Andrews et al., Phys. Rev. Lett. 45, 219(1980).
G. Finnocchiaro et al., Phys. Rev. Lett. 45, 222(1980).

[5] Particle Data Group Review, Phys.Rev.D 54, 1(1996)

[6] Semileptonic Branching Fractions of Charged and Neutral B Mesons
M.Athanas et al., Phys. Rev. Lett. 73, 3503(1994),
(erratum, Phys. Rev. Lett. 74, 3090(1995))

[7] Study of the B^0 Semileptonic Decay Spectrum at the $\Upsilon(4S)$ Resonance
M. Artuso et al., Phys. Lett. B 399, 321(1997).

[8] The Inclusive Decays $B \to DX$ and $B \to D^*X$
L. Gibbons et al., Phys. Rev. D 56, 3783(1997).

[9] Flavor-Specific Inclusive B Decays to Charm
T.E. Coan et al., Phys. Rev. Lett. 80, 1150(1998).

[10] Measurements of $B \to D_s^+ X$ Decays
D. Gibaut et al., Phys. Rev. D 53, 4734(1996).

[11] Inclusive Decays of B Mesons to Charmonium
R. Balest et al., Phys. Rev. D 52, 2661(1995).

[12] The Inclusive Decay $B \to \eta X$
Y. Kubota et al., Phys. Rev. D 53, 6033(1996).

[13] Measurement of the B Semileptonic Branching Fraction with Lepton Tags
B. Barish et al., Phys. Rev. Lett. 76, 1570(1996).

[14] Search for Exclusive Charmless Hadronic B Decays
D.M.Asner et al., Phys. Rev. D 53, 1039(1996).
Evidence for Penguin-Diagram Decays: First Observation of ...
R. Ammar et al., Phys. Rev. Lett. 71, 674(1993).
Observation of $B^+ \to \omega K^+$ and Search for Related B Decay Modes
T. Bergfeld et al., Phys. Rev. Lett. 81, 272(1998).

[15] Measurement of the $\bar{B} \to D\ell\bar{\nu}$ Partial Width and Form Factor Parameters
M. Athanas et al., Phys. Rev. Lett. 79, 2208(1997).

[16] Measurement of the $\bar{B} \to D^*\ell\bar{\nu}$ Branching Fractions and $|V_{cb}|$
B. Barish et al., Phys. Rev. D 51, 1014(1995).

[17] Investigation of Semileptonic B Meson Decay to P-Wave Charm Mesons
A. Anastassov et al., Phys. Rev. Lett. 80, 4127(1998).

[18] First Measurement of the $B \to \pi\ell\nu$ and $B \to \rho(\omega)\ell\nu$ Branching Fractions
J. Alexander et al., Phys. Rev. Lett. 77, 5000(1996).

[19] Exclusive Hadronic B Decays to Charm and Charmonium Final States
M.S. Alam et al., Phys. Rev. D 50, 43(1994).

[20] New Measurement of $B \to D^*\pi$ Branching Fractions
G.Brandenburg et al., Phys. Rev. Lett. 80, 2762(1998).

[21] Measurements of $B \to D_s^+ X$ Decays
D. Gibaut et al., Phys. Rev. D 53, 4734(1996).

[22] Search for the Decays $B^0 \to D^{(*)+}D^{(*)-}$
D.M. Asner et al., Phys. Rev. Lett. 79, 799(1997).

[23] Measurement of the Decay Amplitudes and Branching Fractions of $B \to J/\psi K^*$ and $B \to J/\psi K$ Decays
C.P. Jessop et al., Phys. Rev. Lett. 79, 4533(1997).

[24] Study of $B \to \psi\rho$
M. Bishai et al., Phys. Lett. B 369, 186(1996).

[25] B Meson Decay, B. Gittelman, S. Stone
High Energy Electron-Positron Physics, World Scientific (Singapore), (1988)

[26] Recently there has been a measurement of the branching fraction, $\mathcal{B}r(B \to D^{*-}D^{*+}) = (7.8^{+5.4}_{-3.8} \pm 1.5) \times 10^{-4}$. This has been submitted to the Van Couver conference, CLEO CONF 98-07; ICHEP98 849.



FIG. 1 Side View of the CLEO-II Detector



FIG. 2 The Electro-Weak-Theory Diagrams used to Describe the Most Likely Decays of the B Meson