Diffractive Higgs boson production at LHC

Victor T. Kim
St. Petersburg Nuclear Physics Institute, Gatchina 188300, Russia

Abstract

Predictions for diffractive Higgs boson production at LHC energies are briefly reviewed.

1 Introduction

The advent of high-energy particle colliders such as DESY HERA, Fermilab Tevatron and forthcoming CERN LHC provides a good opportunity to study diffractive (rapidity gap) physics which is intimately related with hard QCD diffraction, i.e., diffraction based on the partonic structure of Pomeron [1]. The partonic structure of Pomeron is established by diffractive jet production in hadron-hadron [2, 3, 4] and lepton-hadron [5] processes.

For intermediate mass Higgs boson ($m_H \leq m_W$) searches at LHC the diffractive processes can be the most important tool for its discovery. The experimental signatures for the heavier Higgs boson mass should be very pronounced, while for the intermediate mass the usual inelastic inclusive Higgs boson production faces a huge background.

Originally, large rapidity gap physics for Higgs boson searches in hadronic collisions via electroweak boson fusion was suggested in Ref. [6]. The idea was that due to colourless exchange by electroweak bosons there is no hadron activity in wide rapidity intervals. Basically one should see Higgs boson signal with rapidity gaps without strong hadronic background.

The double-diffractive Higgs boson production in central region via double-Pomeron exchange, an another process with colourless flow in t-channel and a larger corss section, was proposed in papers [7, 8, 9]. Such a process, shown in Fig. 1, which called also double-Pomeron-exchange process, belongs to the class of multi-Regge processes. The multi-Regge processes, in particular the double-Pomeron exchange, were introduced in Ref. [10] and they have been considered in numerous papers, see, e. g., Refs. [11].

Unfortunately, the available theoretical predictions on double-diffractive Higgs boson production in central region [7], [12]-[18] are very spread in the predicted values for the cross sections. The spread of the predictions is about 10^3(!). This is because of very complicated problems of QCD diffraction, such as factorisation, survival probabilities, etc.

In such circumstances it is impossible to figure out definite expectations for double-diffractive Higgs boson production, which has a unique discovery potential at the Tevatron and especially at the LHC (see Refs. [19]-[21]).

The most important issue concerning the uncertainty is a large factorisation breaking of hard diffractive processes. It is known from data that, according expections of partonic model for Pomeron by Ingelman-Schlein, the diffractive structure function factorises into flux factor of Pomeron and Pomeron structure function in both $\overline{p}p$- and ep- data. However, diffractive structure function extracted from $\overline{p}p$-collisions of CDF at Tevatron is below the ep data from HERA by order of magnitude.

The difference can be caused by the different survival probabilities [22, 23], related with nonperturbative multi-Pomeron exchanges, in hadron-hadron and lepton-hadron processes.

So, now it is generally accepted that the most reliable predictions for diffractive Higgs boson production should reproduce hard difraction data in $\overline{p}p$-collisions of CDF at Tevatron.



Figure 1: The double-diffractive processes for production in central region via double-Pomeron exchange: (a) double-Pomeron exchange for inelastic central production; (b) exclusive central Higgs boson production; (c) inclusive central inelastic Higgs boson production

The other existing $\overline{p}p$-collision data from $S\overline{p}pS$-collider at CERN by UA8 Collaboration are not suitable for complete analysis, because they are uncorrected for detector acceptance/efficiency. Although, soon some new results from UA8 will be available [24].

In the next sections we consider the available predictions for Higgs boson production in double-Pomeron-exchange and single-Pomeron-exchange processes, and present some summary remarks.

2 Higgs boson production in double-Pomeron-exchange processes

Single diffraction cross section via single-Pomeron exchange reads as

$$\frac{d\sigma^{SPE}}{dtd\xi} = F_{I\!P/N}(t,\xi)\cdot\sigma^{tot}_{I\!PN}(s'), \tag{1}$$

$$\tag{2}$$

where ξ is a fraction of the momentum loss by the diffracting nucleon, $-t$ is squared transferred momentum, $\alpha_{I\!P}(t)$ is the Pomeron trajectory, $F_{I\!P/N}(t,\xi)$ is the flux factor of Pomeron.

In the case of the double-diffraction due to double-Pomeron-exchange process with production in central region the following factorisation relation holds (it is obtained by the UA8 Collaboration [3]):

$$\sigma^{tot}_{I\!PI\!P}(s') = \frac{\left[\sigma^{tot}_{I\!PN}(s')\right]^2}{\sigma^{tot}_{NN}(s')}, \tag{3}$$

where $\sigma^{tot}_{I\!P I\!P}$, $\sigma^{tot}_{I\!P N}$ and σ^{tot}_{NN} are the total cross sections of Pomeron-Pomeron, Pomeron-proton and proton-proton collisions, correspondingly.

Therefore, the double-diffraction cross section with production in central region via double-Pomeron exchange, depicted in Fig. 1(a), is of the form:

$$\frac{d\sigma^{DPE}}{d\xi_1 d\xi_2 dt_1 dt_2 d\phi_1 d\phi_2} = F_{I\!P/N}(t_1,\xi_1)\cdot F_{I\!P/N}(t_2,\xi_2)\cdot \sigma^{tot}_{I\!P I\!P}(s'), \tag{4}$$

where ξ_1 and ξ_2 are momentum fractions carried by the collding Pomerons, $s' = \xi_1\xi_2 s = m_X^2$ is the squared invariant mass of the produced system by Pomeron-Pomeron collision.

The hard diffraction, according Ingelman-Schlein, implies that the above Pomeron-proton $\sigma^{tot}_{I\!P N}(s')$ and Pomeron-Pomeron $\sigma^{tot}_{I\!P I\!P}(s')$ total cross sections factorises into parton-parton cross section and Pomeron structure function(s).

The parton distribution of the Pomeron can be taken in the following form:

$$F_{p/I\!P}(x) = (1-P_g)\cdot\Big(F_{q/I\!P}(x) + F_{\overline{q}/I\!P}(x)\Big) + P_g\cdot F_{g/I\!P}(x), \tag{5}$$

where P_g and $(1-P_g)$ are fractions of gluon and quark-antiquark contents of the Pomeron respectively. According diffraction HERA data the partonic structure of the Pomeron is dominant by gluons: $P_g \simeq 0.7 - 0.8$.

If a partonic nature of Pomeron is assumed, then the parton distributions of the effective Pomeron should obey the following momentum sum rule:

$$\begin{aligned}\int_0^1 dx\cdot xF_{N/I\!P}(x) &= \int_0^1 dx\cdot x(2F_{q/I\!P}(x) + F_{g/I\!P}(x)) \\ &= 2<x_q> + <x_g> = (1-P_g) + P_g = 1.\end{aligned} \tag{6}$$

The dominant contribution for diffractive jet (and Higgs) production comes, like in usual inelastic collisions, from gluon distributions. As shown by UA8 data on diffractive jet production [2] gluon distribution in Pomeron can be parameterised in the following way:

$$F_{g/I\!P}(x) = 0.13\cdot 6(1-x)^5/x + 0.57\cdot 6(1-x) + 0.3\cdot\delta(1-x). \tag{7}$$

This parameterisation is compatible with HERA data [5]. Note, that probability about 0.3 to have the "super-hard" gluon component of the Pomeron [2] corresponds to the case when the total momentum of the colliding Pomeron carries by one gluon. A more detail discussion on the "super-hard" Pomeron component as a coherent effect can be found in Refs. [25].

Therefore, the diffractive gluon structure function in nucleon reads as follows:

$$xF^D_{g/N} = xF_{I\!P/N}\otimes F_{g/I\!P}, \tag{8}$$

where $\otimes$ denotes the Mellin convolution of two functions: $f\otimes h \equiv \int_0^1 dy\int_0^1 dz f(y)h(z)\delta(x-yz) = \int_x^1 dy/y f(y)h(x/y)$.

Then the total double-diffractive Higgs boson production cross section in central region is

$$\sigma_H^{DPE} = \int_{m_X} dm \int dx_1 \int dx_2\; F^D_{g/N}(x_1)F^D_{g/N}(x_2)\cdot\sigma_{gg\to H}(M_H, \sqrt{x_1x_2s} = m). \tag{9}$$

An upper estimate of the exclusive double-diffractive Higgs boson production one can make via the "super-hard" component of the Pomeron (Fig. 1(b)).

We present in Fig. 2 the results based on UA8 data at $\sqrt{s} = 630$ [3] for Higgs boson production via double-Pomeron exchange at Tevatron and LHC.

At Tevatron energy $\sqrt{s} = 2$ TeV the predictions yield too low values of the cross section in the exclusive channel to be observed.



Figure 2: The cross section of exclusive $pp \to pHp$ and inclusive $pp \to pHXp$ double-diffractive Higgs boson production suggested by UA8 data for: (a) the Tevatron and (b) the LHC

Even for LHC energy the exclusive Higgs boson production via double-Pomeron-exchange process is practically out of reach: few events per year.

The lowest order perturbative QCD calculation for exclusive double-diffractive Higgs boson production [12] yieds a very tiny cross section: $\sim 10^{-7}$ Fb for $M_H = 120$ GeV at LHC. However, soft contributions and coherence effects drastically increase that result.

There are also predictions [27] based on HERA data on diffractive structure functions. However, they are also very uncertain (the predictions are differ up to factor 600). The problem of compatibility of diffractive cross sections in deep inelastic lepton-hadron and hadron-hadron processes, due to the large factorisation violation, causes a difficulty for reliability of theoretical predictions.

A detail discussion concerning the comparison of those phenomenological calculations one can find in Ref. [21]. According to Ref. [21], the most of the predictions that yield large values for double-diffractive Higgs boson cross sections (≥ 100 Fb) for the LHC will face a problem of considerable overestimating for diffractive jet production at Tevatron [4].

A comparison of the predictions [26], based on UA8 data and available theoretical predictions, is shown in Fig. 3 for the Tevatron and Fig. 4 for the LHC.

3 Higgs production in single-Pomeron-exchange processes

In the original papers [7, 8] for Higgs production in diffractive processes along with the double-Pomeron exchange also the single-Pomeron-exchange mechanism has been proposed (Fig. 5(a)).

It was shown in Ref. [28] that the inclusive single-Pomeron-exchange Higgs boson production is remarkably located in central region like double-Pomeron Higgs production. So, because the single-Pomeron cross section is much larger, one would focus on the inclusive single-Pomeron-exchange Higgs boson production. However, in Refs. [27, 28] is shown that the inclusive single-Pomeron-exchange Higgs boson production does not have a particular advantage at the LHC, though in can be used as an additional channel for double-diffractive Higgs boson production studies. This is due to the fact, that there is a huge backround by three order of magnitude for dominant decay channel with b-quarks for inclusive case.



Figure 3: Comparison of the predictions for the Tevatron, based on UA8 data, with available calculations for: (a) exclusive and (b) inclusive cross sections



Figure 4: Comparison of the predictions for the LHC, based on UA8 data, with available calculations for: (a) exclusive and (b) inclusive cross sections

In case of double-Pomeron exchange, one should use *exclusive* channel, because one could apply final state selection rules [20], which reduce the background by factor about 3000! However, as it was mentioned above the exclusive double-Pomeron-exchange process for Higgs boson productions yields very low rate.

To avoid such problems, in Ref. [29] it has been suggested to consider *exclusive* single-Pomeron-exchange process (see Fig. 5(b) and Fig. 6). The exclusive single-Pomeron-exchange process would be a very important channel to searches of Standard Model Higgs boson with intermediate mass, due to its possibly larger cross section. The caveat of this approach is that the prediction involves rather uncertain factor, related with Higgs boson coupling to proton. In addition, in the exclusive process kinematics the Higgs boson is produced in very forward direction. Experimentally, it requires some special dedicated subdetectors to the existing CMS and ATLAS detectors at the LHC.

4 Summary

The present estimates for exclusive diffractive production of Higgs boson in double-Pomeron-exchange process yield low rates, so it makes rather obscure the outlooks for diffractive Higgs



Figure 5: Single-Pomeron-exchange diagrams for Higgs boson production in: (a) inclusive and (b) exclusive processes



Figure 6: Diagrams for Higgs boson emission (a) after and (b) before the exchange by Pomeron

boson search at LHC (about few events per year).

The new exclusive channel for diffractive Higgs boson production in single-Pomeron-exchange can be very promising, in spite of unusual kinematics of very forward Higgs boson production. Unfortunately, the cross section contains some not very well known parameters which make prediction less certain.

It should be noted, that the most predictions are based on diffractive structure function data from $\overline{p}p$-collisions of CDF at Tevatron, which are below the *ep* data from HERA by order of magnitude. However, preliminary analysis [24] of CERN S$\overline{p}p$S data by UA8 does not show a large difference between $\overline{p}p$- and *ep*- diffractive data. If confirmed, the absence of large factorization breaking can be very important for searches of diffractive Higgs production at Tevatron and LHC colliders. Then Higgs boson production cross section in diffractive processes will be much larger than the current expectations.

References

[1] G. Ingelman and P. E. Schlein, Phys. Lett. B **152** (1985) 256.

[2] R. Bonino *et al.* [UA8 Collaboration], Phys. Lett. B **211** (1988) 239; A. Brandt *et al.* [UA8 Collaboration], Phys. Lett. B **297** (1992) 417; A. Brandt *et al.* [UA8 Collaboration], Nucl. Phys. B **514** (1998) 3 [arXiv:hep-ex/9710004]. A. Brandt *et al.* [UA8 Collaboration], Phys. Lett. B **421** (1998) 395 [arXiv:hep-ex/9709015].



Figure 7: The predictions for inclusive single-Pomeron exchange cross section at Tevatron and LHC energies based on UA8 data.

[3] A. Brandt *et al.* [UA8 Collaboration], Eur. Phys. J. C **25** (2002) 361 [arXiv:hep-ex/0205037].

[4] T. Affolder *et al.* [CDF Collaboration], Phys. Rev. Lett. **85** (2000) 4215; B. Abbott *et al.* [D0 Collaboration], Phys. Lett. B **531** (2002) 52 [arXiv:hep-ex/9912061].

[5] T. Ahmed *et al.* [H1 Collaboration], Phys. Lett. B **348** (1995) 681 [arXiv:hep-ex/9503005]. M. Derrick *et al.* [ZEUS Collaboration], Z. Phys. C **68** (1995) 569 [arXiv:hep-ex/9505010].

[6] Yu. L. Dokshitzer, V. A. Khoze and S. I. Troyan, LNPI-86-1218, Leningrad (1986), *Proc. Physics in Collision VI, Chicago, USA, 1986, ed. M. Derrick (World Scientific, Singapore, 1987) pp. 417-448*; Sov. J. Nucl. Phys. **46** (1987) 712 [Yad. Fiz. **46** (1987) 1220]; Yu. L. Dokshitzer, V. A. Khoze and T. Sjostrand, Phys. Lett. B **274** (1992) 116.

[7] A. Schäfer, O. Nachtmann and R. Schöpf, Phys. Lett. B **249** (1990) 331; A. Białas and P. V. Landshoff, Phys. Lett. B **256** (1991) 540.

[8] B. Müller and A. J. Schramm, Nucl. Phys. A **523** (1991) 677.

[9] J. D. Bjorken, Phys. Rev. D **47** (1993) 101.

[10] K. A. Ter-Martirosyan, Zh. Eksp. Teor. Fiz. **44** (1963) 341 [Sov. Phys. JETP **17** (1963) 223]; T. W. Kibble, Phys. Rev. **131** (1963) 2282; Z. Koba, Fortsch. Phys. **11** (1963) 118.

[11] R. Shankar, Nucl. Phys. B **70** (1974) 168; D. M. Chew, Nucl. Phys. B **82** (1974) 422; D. M. Chew and G. F. Chew, Phys. Lett. B **53** (1974) 191; A. B. Kaidalov and K. A. Ter-Martirosyan, Nucl. Phys. B **75** (1974) 471; Ya. I. Azimov, V. A. Khoze, E. M. Levin and M. G. Ryskin, Yad. Fiz. **21** (1975) 413 [Sov. J. Nucl. Phys. **21** (1975) 215]; B. R. Desai, B. C. Shen and M. Jacob, Nucl. Phys. B **142** (1978) 258.

[12] H. J. Lu and J. Milana, Phys. Rev. D **51** (1995) 6107 [arXiv:hep-ph/9407206].

[13] J.-R. Cudell and O. F. Hernandez, Nucl. Phys. B **471** (1996) 471 [arXiv:hep-ph/9511252].

[14] V. A. Khoze, A. D. Martin and M. G. Ryskin, Phys. Lett. B **401** (1997) 330 [arXiv:hep-ph/9701419]; Eur. Phys. J. C **14** (2000) 525 [arXiv:hep-ph/0002072]; Eur. Phys. J. C **19** (2001) 477 [Erratum-ibid. C **20** (2001) 599] [arXiv:hep-ph/0011393].

[15] D. Kharzeev and E. Levin, Phys. Rev. D **63** (2001) 073004 [arXiv:hep-ph/0005311]; E. Levin, arXiv:hep-ph/9912403.

[16] M. Boonekamp, R. Peschanski and C. Royon, Phys. Rev. Lett. **87** (2001) 251806 [arXiv:hep-ph/0107113]; M. Boonekamp, A. De Roeck, R. Peschanski and C. Royon, arXiv:hep-ph/0205332.

[17] B. Cox, J. Forshaw and B. Heinemann, Phys. Lett. B **540** (2002) 263 [arXiv:hep-ph/0110173].

[18] R. Enberg, G. Ingelman, A. Kissavos and N. Timneanu, Phys. Rev. Lett. **89** (2002) 081801 [arXiv:hep-ph/0203267].

[19] M. G. Albrow and A. Rostovtsev, Fermilab-Pub-00-173 (2000), arXiv:hep-ph/0009336.

[20] A. De Roeck, V. A. Khoze, A. D. Martin, R. Orava and M. G. Ryskin, Eur. Phys. J. C **25** (2002) 391 [arXiv:hep-ph/0207042].

[21] V. A. Khoze, A. D. Martin and M. G. Ryskin, IPPP-02-44, DCPT-02-88 (2002), arXiv:hep-ph/0207313.

[22] E. Gotsman, E. Levin and U. Maor, Phys. Rev. D **60** (1999) 094011 [arXiv:hep-ph/9902294]; Phys. Lett. B **438** (1998) 229 [arXiv:hep-ph/9804404].

[23] A. B. Kaidalov, V. A. Khoze, A. D. Martin and M. G. Ryskin, Eur. Phys. J. C **21** (2001) 521 [arXiv:hep-ph/0105145].

[24] V. T. Kim and P. E. Schlein, "Absence of large factorization breaking in hard diffractive data," in preparation.

[25] J. C. Collins, L. Frankfurt and M. Strikman, Phys. Lett. B **307** (1993) 161 [arXiv:hep-ph/9212212]; J. C. Collins, J. Huston, J. Pumplin, H. Weerts and J. J. Whitmore, Phys. Rev. D **51** (1995) 3182 [arXiv:hep-ph/9406255].

[26] S. Erhan, V. T. Kim and P. E. Schlein, 38th PNPI Winter School Repino, February 2003.

[27] D. Graudenz and G. Veneziano, Phys. Lett. B **365** (1996) 302 [arXiv:hep-ph/9508401]; M. Heyssler, Z. Kunszt and W. J. Stirling, Phys. Lett. B **406** (1997) 95 [arXiv:hep-ph/9702286].

[28] S. Erhan, V. T. Kim and P. E. Schlein, CERN-TH-2003-232, UCLA-EPP-2003-101, arXiv:hep-ph/0312342.

[29] V. T. Kim, IX RDMS CMS Conference, Minsk, November 2004;
X RDMS CMS Conference, St. Petersburg, September 2005.