

Shadows, currents and AdS

R.R. Metsaev

Lebedev Institute

hep-th/

Plan

- 1) gauge invariant formulation of conformal currents and shadow fields
- 2) AdS/CFT correspondence for these currents and shadows and massless AdS fields

Spin-1 conformal current

Standard approach

T^a — *conformal current*

$$\partial^a T_a = 0$$

Conformal dimension

$$\Delta = d - 1$$

Spin-1 conformal current. Gauge inv. approach

$$T^a = \phi_{\text{cur}}^a + \partial^a \phi_{\text{cur}}$$

$$\partial^a T^a = 0$$

$$\partial^a \phi_{\text{cur}}^a + \square \phi_{\text{cur}} = 0$$

$$\delta \phi_{\text{cur}}^a = \partial^a \xi$$

$$\delta \phi_{\text{cur}} = -\xi$$

Conformal dimensions

$$\Delta(T^a) = d - 1$$

$$T^a = \phi_{cur}^a + \partial^a \phi_{cur}$$

$$\Delta(\phi_{cur}^a) = d - 1$$

$$\Delta(\phi_{cur}) = d - 2$$

Shadow spin-1 field: Definition

T^a *current*

Φ^a *is shadow field IF*

$$\mathcal{L} = \Phi^a T^a$$

is invariant w.r.t conformal

symmetries $so(d,2)$

$$\delta\Phi^a = \partial^a \xi$$

Spin-1 shadow field. Gauge inv. approach

$$\phi_{\text{sh}}^{\mathbf{a}} \quad \phi_{\text{sh}}$$

$$\partial^{\mathbf{a}} \phi_{\text{sh}}^{\mathbf{a}} + \phi_{\text{sh}} = 0$$

$$\delta \phi_{sh}^a = \partial^a \xi_{sh}$$

$$\delta \phi_{sh} = -\square \xi_{sh}$$

Conformal dims. Shadows

$$\Delta(\phi^a) = 1$$

$$\partial^a \phi_{sh}^a + \phi_{sh} = 0$$

$$\Delta(\phi_{sh}^a) = 1$$

$$\Delta(\phi_{sh}) = 2$$

Step 2

One needs to prove that
gauge invariant approach respects
conformal symmetries
of $so(d, 2)$ algebra

Conformal algebra $so(d, 2)$

P^a translations

J^{ab} Lorentz rotations

D Dilatation

K^a conformal boosts

**All that remains is to respect
conformal boost symmetries**

K^a

$$K^a = \tilde{K}^a + \mathbf{R}^a$$

$$\tilde{K}^a \equiv -\frac{1}{2}x^2\partial^a + x^a D + M^{ab}x^b$$

we have to find operator

$$R^a$$

Currents

$$R^a \phi_{cur}^b = \eta^{ab} \phi_{cur}$$

$$R^a \phi_{cur} = 0$$

Shadows

$$R^a \phi_{sh}^b = 0$$

$$R^a \phi_{sh} = \phi_{sh}^a$$

Spin-2 current. Gauge inv. approach

Fields

Conf.dim

$\phi_{\text{cur}}^{\text{ab}}$

d

$\phi_{\text{cur}}^{\text{a}}$

$d - 1$

ϕ_{cur}

$d - 2$

Spin 2. Currents.

Differential constraints

$$\partial^b \phi_{cur}^{ab} + \partial^a \phi_{cur}^{bb} + \square \phi_{cur}^a = 0$$

$$\partial^a \phi_{cur}^a + \phi_{cur}^{aa} + \square \phi_{cur} = 0$$

ϕ_{cur}^a , ϕ_{cur} can be gauged away

$$\partial^b \phi_{cur}^{ab} = 0$$

$$\phi_{cur}^{aa} = 0$$

Spin 2. Currents.

Gauge transformations

$$\delta\phi_{cur}^{ab} = \partial^a \xi_{cur}^b + \partial^b \xi_{cur}^a + \eta^{ab} \square \xi_{cur}$$

$$\delta\phi_{cur}^a = \partial^a \xi_{cur} + \xi_{cur}^a$$

$$\delta\phi_{cur} = \xi_{cur}$$

ϕ_{cur}^a, ϕ_{cur} Stueckelberg fields

Spin 2. Currents.

Gauge fields and energy-momentum tensor

$$T^{ab} = \phi_{cur}^{ab} + \partial^a \phi_{cur}^b + \partial^b \phi_{cur}^a + \partial^a \partial^b \phi_{cur} + \eta^{ab} \square \phi_{cur}$$

1) T^{ab} is gauge invariant

$$2) \partial^a T^{ab} = 0, \quad T^{aa} = 0$$

amount to differential constraints
for ϕ_{cur}^{ab} , ϕ_{cur}^a , ϕ_{cur} .

Spin 2. Shadows

Fields

Conf.dim

$\phi_{\text{sh}}^{\text{ab}}$

0

$\phi_{\text{sh}}^{\text{a}}$

1

ϕ_{sh}

2

Spin 2. Shadows

differential constraints

$$\partial^b \phi_{sh}^{ab} + \partial^a \phi_{sh}^{bb} + \phi_{sh}^a = 0$$

$$\partial^a \phi_{sh}^a + \square \phi_{sh}^{aa} + \phi_{sh} = 0$$

ϕ_{sh}^{aa} can be gauged away

$$\partial^b \phi_{sh}^{ab} + \phi_{sh}^a = 0$$

$$\partial^a \phi_{sh}^a + \phi_{sh} = 0$$

Spin 2. Shadows gauge transformations

$$\delta\phi_{sh}^{ab} = \partial^a \xi_{sh}^b + \partial^b \xi_{sh}^a + \eta^{ab} \xi_{sh}$$

$$\delta\phi_{sh}^a = \partial^a \xi_{sh} + \square \xi_{sh}^a$$

$$\delta\phi_{sh} = \square \xi_{sh}$$

ϕ_{sh}^{aa} Stueckelberg field

ϕ_{sh}^a, ϕ_{sh} are not Stueckelberg fields

Generalization to

arbitrary spin currents and shadows

is straightforward by using

double-traceless

Fronsdal fields

Statement

Requiring

1) gauge invariance

2) conformal invariance $so(d,2)$

we obtain solution to

differential constraints

and operator R^a

AdS/CFT

Our currents and shadows

correspond to

bulk AdS fields taken

in modified de Donder gauge

summary of our study of AdS/CFT

1) Bulk fields are taken in

modified de-Donder gauge

2) **de Donder gauge** leads to

decoupled equations of motion

with on-shell leftover gauge symmetries

3) normalizable solutions \rightarrow currents

no-normalizable solutions \rightarrow shadows

4) leftover on-shell gauge symmetries of bulk fields correspond to off-shell gauge symmetries of boundary currents and shadow fields,

5) **de Donder gauge** for bulk fields corresponds to **differential constraints** for boundary conformal currents and shadows

AdS/CFT

AdS_{d+1} space

$$ds^2 = \frac{1}{z^2}(dx^a dx^a + dz dz)$$

bulk $so(d, 1) \rightarrow$ **boundary** $so(d - 1, 1)$

$$\phi^A = \phi^a \oplus \phi$$

AdS. Spin 1

$$D^A F^{AB} = 0$$

$$F^{AB} = D^A \phi^B - D^B \phi^A$$

$$\phi^A = \phi^{\mathbf{a}} \oplus \phi$$

(in radial gauge, $\phi = 0$)

standard Lorentz gauge

$$D^A \phi^A = 0$$

leads to

coupled equations

$$(\square + \partial_z^2 - m^2) \phi^{\mathbf{a}} + \partial^a \phi = 0$$

$$(\square + \partial_z^2 - m^2) \phi + \partial^a \phi^{\mathbf{a}} = 0$$

Modified Lorentz gauge

$$D^A \phi^A + 2\phi = 0$$

RRM, 1999

Polchinski and

Strassler 2001

gives

Decoupled equations

Decoupled equations

$$(\square + \partial_z^2 - m_1^2)\phi^a = 0$$

$$(\square + \partial_z^2 - m_0^2)\phi = 0$$

$$m_1^2 = \frac{1}{z^2}(\nu_1^2 - \frac{1}{4})$$

$$m_0^2 = \frac{1}{z^2}(\nu_0^2 - \frac{1}{4})$$

$$\phi^a(x,z) = \mathbf{U}_{\nu_1} \phi^a_{cur}(x)$$

$$\phi(x,z) = \mathbf{U}_{\nu_0} \phi_{cur}(x)$$

$$\mathbf{U}_{\nu} \equiv \sqrt{z} \mathbf{J}_{\nu}(z\sqrt{\square})$$

Bessel

modified Lorentz gauge

$$\partial^{\mathbf{a}}\phi^{\mathbf{a}} + \left(\partial_{\mathbf{z}} + \frac{a}{z}\right)\phi = 0$$

$$\partial^{\mathbf{a}}\phi^{\mathbf{a}} + \left(\partial_{\mathbf{z}} + \frac{a}{z}\right)\phi$$

$$= U_{\nu_1}(\partial^{\mathbf{a}}\phi^{\mathbf{a}}_{\text{cur}} + \square\phi_{\text{cur}})$$

**“technical” problem with standard
cov. gauges, Lorentz, de Donder**

1) Coupled equations

2) For spin 2, 3, 4,

solutions are expressible

in terms of **Heun functions**

Little is known about **Heun functions**

asymptotic behavior ???

recurrent relations ???

Spin 2: modified de Donder gauge

$$D^B h^{AB} - \frac{1}{2} D^A h + 2h^{zA} - \eta^{zA} h = 0$$

$$h \equiv h^{AA}$$

leads to **decoupled** equations

$$h^{AB} = h^{ab} \oplus h^{za} \oplus h^{zz}$$

$$T^{ab} = \phi_{cur}^{ab} + \partial^a \phi_{cur}^b + \partial^b \phi_{cur}^a + \partial^a \partial^b \phi_{cur} + \eta^{ab} \square \phi_{cur}$$

Breaking conformal symmetry

$$\square \rightarrow m^2$$

$$\phi_{cur}^a \rightarrow \frac{1}{m} \phi^a \qquad \phi_{cur} \rightarrow \frac{1}{m^2} \phi^a$$

$$T^{ab} = \phi^{ab} + \frac{1}{m} (\partial^a \phi^b + \partial^b \phi^a) + \frac{\partial^a \partial^b}{m^2} \phi + \eta^{ab} \phi$$

Conclusions

1) Gauge invariant approach to currents and shadows

give possibility to choice

various gauges which might be helpful in applications

Standard currents are obtained via Stueckelberg gauge fixing.

But others gauges might also be interesting

2)

**modified de Donder gauge leads to
decoupled equations**

and might be helpful for study

AdS/CFT

AdS/QCD

quantization of higher-spin

AdS fields