

Asymptotic effects in jet production at high energies*

V.B. Gavrilov[&], V.T. Kim[‡], A.A. Krokhotin[&], G.B. Pivovarov[‡] and G.B. Safronov[&]

& Institute of Theoretical and Experimental Physics, Moscow, 117218 Russia

‡ St. Petersburg Nuclear Physics Institute, Gatchina, 188300 Russia

‡ *Institute for Nuclear Research, Moscow, 117312 Russia*

Abstract

Monte Carlo event simulation with BFKL evolution, which governs the high-energy asymptotics of inclusive cross sections, is discussed. We report current status of a Monte Carlo event generator ULYSSES with implemented BFKL evolution. The ULYSSES, based on Pythia Monte Carlo generator, would help to reveal BFKL effects at LHC energies. In particular, such an observable as dijet K-factor can serve as a source of BFKL dynamics at the LHC, and it also would help for search of new physics.

With advent of the Large Hadron Collider (LHC) at CERN we have an opportunity to probe the Standard Model far beyond the explored domains. QCD is an essential ingredient of the Standard Model, and it is well tested in hard processes when transferred momentum is of the order of the total collision energy (Bjorken limit: $Q^2 \sim s \rightarrow \infty$). The cornerstones of perturbative QCD at this kinematic regime (QCD-improved parton model): factorization of inclusive hard processes [1] and the Gribov–Lipatov–Altarelli–Parisi–Dokshitzer (GLAPD) evolution equation [2] provides a basis for the successful QCD-improved parton model. The factorization theorem [1] for inclusive hard processes ensures that the inclusive cross section factorizes into partonic subprocess(es) and parton distribution function(s). The GLAPD-evolution equation governs the $\log Q^2$ -dependence (at $Q^2 \rightarrow \infty$) of the inclusive hard process cross-sections at fixed scaling variable $x \sim Q^2/s$.

Another kinematic domain that is very important at high-energy is given by the Balitsky–Fadin–Kuraev–Lipatov (BFKL) limit [3, 4, 5, 6], or QCD-Regge limit, whereby at fixed $Q^2 \gg \Lambda_{QCD}^2$, $s \rightarrow \infty$. In the BFKL limit, the BFKL evolution in the leading-log approximation (LLA) governs $\log(1/x)$ evolution (at $x \rightarrow 0$) of inclusive processes.

One of the key BFKL feature is relation of the highest eigenvalue, ω^{max} , of the BFKL equation [3, 4, 5, 6] with the intercept of the Pomeron, which in turn governs the high-energy asymptotics of the total cross-sections: $\sigma \sim (s/s_0)^{\alpha_{IP}-1} = (s/s_0)^{\omega^{max}}$, where the Regge parameter s_0 defines the approach to the asymptotic regime. The BFKL Pomeron intercept in the LLA turns out to be rather large: $\alpha_{IP} - 1 = \omega_{LLA}^{max} = 12 \log 2 (\alpha_S/\pi) \simeq 0.54$ for $\alpha_S = 0.2$; hence, it is very important to take into account NLLA corrections [7, 8, 9, 11, 10] to the BFKL. Note that the BFKL evolution in the next-to-leading-log approximation (NLLA) [7, 8, 12], unlike the LLA BFKL [3, 4, 5], includes GLAPD evolution with the running coupling constant of the leading-order (LO) GLAPD, $\alpha_S(Q^2) = 4\pi/\beta_0 \log(Q^2/\Lambda_{QCD}^2)$.

One of the striking features of the NLLA BFKL analysis [8] is that the NLLA value for the intercept of the BFKL Pomeron, improved by the BLM procedure [13], has a very weak dependence on the gluon virtuality Q^2 : $\alpha_{IP} - 1 = \omega_{NLLA}^{max} \simeq 0.13 - 0.18$ at $Q^2 = 1 - 100 \text{ GeV}^2$. This agrees with the conventional Regge theory where one expects universal intercept of the Pomeron without any Q^2 -dependence. The value of NNLA BFKL Pomeron intercept [8] becomes compatible with the available data on hard QCD Pomeron. So, NLLA BFKL approach

*presented by V.T.K. at QUARKS'2008, Zagorsk, Russia, May 25-31, 2008

[8, 12] possess all basic features of BFKL, but include also running coupling effects and realistic values of the hard Pomeron intercept.

Therefore, the BFKL and especially the NLLA BFKL [7, 8, 12] are anticipated to be important tools for exploring the high-energy limit of QCD.

It should be stressed that in contrast to the GLAPD, BFKL dynamics involves unintegrated over k_t parton distributions. So, to reveal BFKL effects one needs to deal with, e.g., with parton (jet) production. For jet production in the LLA BFKL within k_t -factorization [14] one can use effective Feynman-like rules for inclusive jet cross sections [15].

However, extremely sophisticated design of contemporary high-energy detectors require from theory and phenomenology a Monte Carlo event generator implementation. Widely used Monte Carlo event generator Pythia [16] contains basically GLAPD-evolution. To incorporate BFKL-evolution the Pythia has been modified in parton shower kernel and parton distribution functions.

For Monte Carlo event simulations of GLAPD evolution is convenient to use the Dokshitzer-Diakonov-Troyan (DDT) [17] presentation:

$$\frac{\partial a(x, k_t^2)}{\partial \log k_t^2} = \frac{\alpha_S(k_t^2)}{2\pi} \sum_b \int_x^1 dz P_{ab}(z) b\left(\frac{x}{z}, k_t^2\right) - \frac{a(x, k_t^2)}{T_a(k_t^2, \mu^2)} \frac{\partial T_a(k_t^2, \mu^2)}{\partial \log k_t^2},$$

where $P_{ab}(z)$ are kernels (splitting functions) of GLAPD equation for partons a and b ,

$$a(x, \mu^2) = \int^{k_t^2} dk_t^2 f_a(x, k_t^2, \mu^2), \quad a = g, q, \bar{q}$$

is GLAPD parton distribution for parton a , $f_a(x, k_t^2, \mu^2)$ is so-called unintegrated parton distribution used in BFKL approach and

$$T_a(k_t^2, \mu^2) \equiv \exp\left(-\int_{k_t^2}^{\mu^2} dk_t^2 \frac{\alpha_S(k_t^2)}{2\pi} \sum_b \int_0^1 d\zeta \zeta P_{ba}(\zeta)\right)$$

is QCD analog for Sudakov form factor (DDT form factor [17]), which defines the probability not to emit (or to emit) a gluon.

In terms of unintegrated parton distribution the GLAPD equation can be represented as:

$$\begin{aligned} f_a(x, k_t^2, \mu^2) &\equiv \frac{\partial}{\partial \log k_t^2} [a(x, k_t^2) T_a(k_t^2, \mu^2)] \\ &= T_a(k_t^2, \mu^2) \frac{\alpha_S(k_t^2)}{2\pi} \sum_b \int_x^1 dz P_{ab}(z) b\left(\frac{x}{z}, k_t^2\right) \end{aligned}$$

Then, a unified BFKL-GLAPD equation can be presented in the following way [18]:

$$\begin{aligned} f_g(x, k_t^2, \mu^2) &= T_g(k_t, \mu) \frac{\alpha_S(k_t^2)}{2\pi} \left\{ \int_x^{\mu/(\mu+k_t)} dz \int^{k_t^2} \frac{dk_t'^2}{k_t'^2} \left[\bar{P}(z) h_g\left(\frac{x}{z}, k_t'^2\right) \right. \right. \\ &+ \left. \left. P_{gq}(z) \sum h_q\left(\frac{x}{z}, k_t'^2\right) \right] + 2N_C \int_x^{\mu/(\mu+k_t)} \frac{dz}{z} \int \frac{d^2q}{\pi q^2} \right. \\ &\times \left. \left[\frac{k_t^2}{k_t'^2} h_g\left(\frac{x}{z}, k_t'^2\right) - \Theta(k_t^2 - q^2) h_g\left(\frac{x}{z}, k_t^2\right) \right] \right\}, \quad \bar{P}(z) = P_{gg}(z) - \frac{2N_C}{z} \end{aligned} \quad (1)$$

The Eq. (1) is similar to Ciafaloni-Catani-Fiorani-Marchesini (CCFM) [19] equation, which is a version of BFKL equation with imposed angle ordering.




Figure 1: Dijet K-factor at LHC: inclusive dijet cross section over only-two-jet cross section ratio. The results are shown with full CMS detector simulation (arbitrary normalization) for luminosity pb^{-1} .

Monte Carlo event generator Ulysses [20] contains an implementation of the Eq. (1) into Pythia [16] parton shower simulation and embedded BFKL unintegrated parton distributions. Preliminary results for dijet K-factor at CMS detector of the LHC are shown in Fig. (), where dijet K-factor for jets with $E_t > E_t^{min} = 60$ GeV is defined as ratio of inclusive dijet cross section to "exclusive" one. "Exclusive" dijet event means the event when only two jets with E_t above E_t^{min} are available ("Born" cross section).

To summarize, BFKL Monte Carlo should help to look at LHC for BFKL dynamics as a new feature of asymptotic QCD. It also has a potential for search of new physics, e.g., graviton production in trans-Planckian regime [21].

V.T.K. thanks the organizers of the QUARKS'2008 for their warm hospitality. This work was supported in parts by RFBR grant No. 08-02-01184a, and RF President grants Nos. 1616.2008.2 and 378.2008.2.

References

- [1] D. Amati, R. Petronzio and G. Veneziano, Nucl. Phys. **B140**, 54 (1978); *ibid.* **B146**, 29 (1978);
S. B. Libby and G. Sterman, Phys. Rev. **D18**, 3252 (1978);
A. H. Mueller, Phys. Rev. **D18**, 3705 (1978);
A. V. Efremov and A. V. Radyushkin, JINR-E2-11725, 11726, 11849, Dubna (1978),
Teor. Mat. Fiz. **44**, 17, 157, 327 (1980) [J. Theor. Math. Phys. **44**, 573, 664, 774 (1981)];
R. K. Ellis, H. Georgi, M. Machacek, H. D. Politzer and G. G. Ross, Nucl. Phys. **B152**, 285 (1979);
J. C. Collins, D. E. Soper and G. Sterman, Nucl. Phys. **B261**, 104 (1985);
G. T. Bodwin, Phys. Rev. **D31**, 2616 (1985); *ibid.* (E) **D34**, 3932 (1986);
for a review, see J. C. Collins, D. E. Soper and G. Sterman, in: *Perturbative QCD*, ed. A. H. Mueller (World Scientific, Singapore, 1989) p. 1.
- [2] V. N. Gribov, L. N. Lipatov, Yad. Fiz. **15**, 781, 1218 (1972) [Sov. J. Nucl. Phys. **15**, 438, 675 (1972)];
L. N. Lipatov, Yad. Fiz. **20**, 181 (1974) [Sov. J. Nucl. Phys. **20**, 94 (1975)];
G. Altarelli and G. Parisi, Nucl. Phys. **B126**, 298 (1977);
Yu. L. Dokshitzer, ZhETF **73**, 1216 (1977) [Sov. Phys. JETP **46**, 641 (1977)].
- [3] E. A. Kuraev, L. N. Lipatov and V. S. Fadin, Phys. Lett. **B60**, 50 (1975); ZhETF **71**, 840 (1976) [Sov. Phys. JETP **44**, 443 (1976)]; *ibid.* **72**, 377 (1977) [**45**, 199 (1977)];
L. N. Lipatov, Yad. Fiz. **23**, 642 (1976) [Sov. J. Nucl. Phys. **23**, 338 (1976)]; ZhETF **90**, 1536 (1986) [Sov. Phys. JETP **63**, 904 (1986)].
- [4] I. I. Balitsky and L. N. Lipatov, Yad. Fiz. **28**, 1597 (1978) [Sov. J. Nucl. Phys. **28**, 822 (1978)].
- [5] L. N. Lipatov, Phys. Rep. **C286**, 131 (1997).
- [6] L. V. Gribov, E. M. Levin and M. G. Ryskin, Phys. Rep. **C100**, 1 (1983).
- [7] V. S. Fadin and L. N. Lipatov, Phys. Lett. **B429**, 127 (1998);
G. Camici and M. Ciafaloni, Phys. Lett. **B430**, 349 (1998); and references therein.
- [8] S. J. Brodsky, V. S. Fadin, V. T. Kim, L. N. Lipatov and G. B. Pivovarov, Pisma ZhETF **70**, 161 (1999) [JETP Lett. **70**, 155 (1999)], hep-ph/9901229.
- [9] M. Ciafaloni, D. Colferai, and G. P. Salam, Phys. Rev. **D60** (1999) 114036;
M. Ciafaloni and D. Colferai, Phys. Lett. **B452** (1999) 372.
- [10] G. Altarelli, R. D. Ball, and S. Forte, Nucl. Phys. **B575** (2000) 313.
- [11] R. S. Thorne, Phys. Rev. **D60** (1999) 054031; *ibid.* **D64** (2001) 074005.
- [12] V. T. Kim, L. N. Lipatov and G. B. Pivovarov, IITAP-99-013 Ames (1999), hep-ph/9911228; IITAP-99-014 Ames (1999), hep-ph/9911242.
- [13] S. J. Brodsky, G. P. Lepage and P. B. Mackenzie, Phys. Rev. **D28**, 228 (1983).
- [14] S. Catani, M. Ciafaloni and F. Hautmann, Nucl. Phys. **B366**, 135 (1991); J. C. Collins and R. K. Ellis, Nucl. Phys. **B360**, 3 (1991); E. M. Levin, M. G. Ryskin, Yu. M. Shabelsky and A. G. Shuvaev, Yad. Fiz. **54** 1420 (1991) [Sov. J. Nucl. Phys. **54**, 867 (1991)].
- [15] V. T. Kim, G. B. Pivovarov, Phys. Rev. **D54**, 725 (1996).

- [16] T. Sjostrand, S. Mrenna, P. Skands, JHEP **0605**, 026 (2006).
- [17] Yu. L. Dokshitzer, D. I. Diakonov and S. I. Troyan, Phys. Rep. **C92**, 90 (1980);
- [18] M. A. Kimber, J. Kwiecinski, A. D. Martin and A. M. Stasto, Phys.Rev. **D62**, 4006 (2000);
M. A. Kimber, A. D. Martin and M. G. Ryskin, Phys.Rev. **D63**, 114027 (2001).
- [19] M.Ciafaloni, Nucl. Phys. **B296**, 49 (1988);
S. Catani, F. Fiorani, G. Marchesini, Phys. Lett. **B234**, 339 (1990); Nucl. Phys. **B336**, 18 (1990);
G. Marchesini, Nucl. Phys. **B445**, 49 (1995)
- [20] V. T. Kim, A. A. Krokhotin and G. B. Pivovarov, in prepration.
- [21] G. F. Giudice, R. Rattazzi, J. D. Wells, Nucl.Phys. **B544**, 3 (1999); Nucl. Phys. **B630**, 293 (2002).