

# On two-point correlation functions in AdS/QCD

A. A. Krikun<sup>a\*</sup>

<sup>a</sup> *MIPT and ITEP*

*Bolshaya Cheremushkinskaya, 25, Moscow, Russia*

## Abstract

In this talk we study the hard wall AdS/QCD model. The simplest model of bottom-up approach to AdS/QCD. We reveal the relations between fields in the model and operators in QCD, fix parameters of the model and calculate some results. We underline the problems of the model and propose the way to solve them.

Presented at 15th International Seminar “Quarks-2008”, Sergiev Posad, Russia, May 2008

## 1 Introduction

In the last few years a great attention was paid to the so-called phenomenological AdS/QCD theories. The essence of these models is to use the AdS/CFT correspondence [2] to describe QCD in large  $N_c$  limit via its 5-dimensional dual theory. The exact structure of this 5D theory, describing all specific features of QCD is not clear, but some simple models have been proposed [3] [4] [5] [6] [7] [8], which already give promising results.

In this talk we study the simplest of these settings, the so-called hard wall AdS/QCD model (see for example [3] [5], first proposed in [9]). Our goal is to find proper relation between fields in the theory and QCD currents, fix the free parameters of the model and study results, that this model gives without any tuning. After that we propose some ways to tune up the model in order to reproduce results of QCD. We will calculate vector, axial vector and pseudoscalar two-point functions of QCD, that will allow us to find values of meson masses and  $f_\pi$ .

This talk is mainly based on the paper by A.K. [1], so all details and references can be seen there.

## 2 Description of the model

We consider the simplest holographic model of low energy QCD, proposed in [3],[5],[7](see also [14, 15, 16]), the so-called “Hard wall AdS/QCD model”. In what follows we will work with conventions and notations, used in [3].

In the AdS/CFT prescription, the fields in 5-dimensional space are dual to operators in 4D, and the global flavor symmetry of the 4D field theory corresponds to the gauge symmetry in its 5D dual. So we will study 4D QCD with  $SU(2)_L \times SU(2)_R$  global symmetry via the gauge theory in AdS with  $SU(2)_L \times SU(2)_R$  gauge group. In this model only the fields, dual to QCD operators with the lowest dimensions, are considered.

We have the  $SU(2)_L \times SU(2)_R$  gauge field theory in  $AdS_5$  space with the metric:

$$ds^2 = \frac{R^2}{z^2}(-dz^2 + dx^\mu dx_\mu) \quad (1)$$

---

\* e-mail: e-mail: krikun.a@gmail.com

where  $R$  is the AdS curvature radius, cut at  $z$  coordinate:  $0 < z \leq z_m$

Later, we will denote 5-dimensional indices with  $A, B, \dots \in \{0, 1, 2, 3, z\}$ , and 4D indices with  $\mu, \nu, \dots \in \{0, 1, 2, 3\}$ .

The theory includes left- and right-handed gauge vector fields  $SU_L(2) \times SU_R(2)$  ( $A_L$  and  $A_R$ , respectively) and bifundamental scalar  $X_{\alpha\beta}$ . According to AdS/CFT 5D fields correspond to operators in QCD:

$$\begin{aligned} A_{L\mu}^a &\leftrightarrow \bar{q}_L \gamma^\mu t^a q_L \\ A_{R\mu}^a &\leftrightarrow \bar{q}_R \gamma^\mu t^a q_R \\ \left(\frac{2}{z}\right) X^{\alpha\beta} &\leftrightarrow \bar{q}_R^\alpha q_L^\beta \end{aligned}$$

with the boundary conditions imposed at  $z = z_m$ :

$$\partial_z V(z_m) = 0 \quad ; \quad \partial_z A(z_m) = 0.$$

The action is:

$$S = \int d^5x \sqrt{g} Tr \left\{ \Lambda^2 (|DX|^2 + \frac{3}{R^2} |X|^2) - \frac{1}{4g_5^2} (F_L^2 + F_R^2) \right\}$$

where

$$\begin{aligned} D_B X &= \partial_B X - \imath A_{LB} X + \imath X A_{RB} \\ A_{L(R)} &= A_{L(R)}^a t^a \\ F_{BD} &= \partial_B A_D - \partial_D A_B - \imath [A_B, A_D], \end{aligned}$$

and we introduce the normalization constant  $\Lambda$  of field  $X$ .

From the equation of motion for  $X$  we can get classical solution:

$$X_0(z) = \frac{1}{2} M z + \frac{1}{2} \Sigma z^3.$$

According to AdS/CFT [17], we argue, that  $M$  corresponds to quark mass matrix, i.e. the source of operator  $\bar{q}_R^\alpha q_L^\beta$  and  $\Sigma$  to condensates, i.e. vacuum expectation value of  $\bar{q}_R^\alpha q_L^\beta$ . We can make  $M$  to be quark masses exactly by appropriate definition of normalization  $\Lambda$ , but the relation between  $\Sigma$  and condensates is to be ascertained. In further discussion we set  $\Sigma = \sigma \mathbf{1}$ ,  $M = m \mathbf{1}$ , assuming the equality of quark masses.

$$X_0(z) = \frac{1}{2} v(z) \mathbf{1}, \quad v(z) = mz + \sigma z^3$$

We will decompose  $X$  on modulus and phase:

$$X = X_0 e^{\imath 2\pi^a (t^a)} = \mathbf{1} \frac{v(z)}{2} e^{\imath 2\pi^a t^a}$$

It is convenient to introduce the vector and axial vector fields:

$$\begin{aligned} V &= (A_L + A_R)/2 \\ A &= (A_L - A_R)/2. \end{aligned}$$

We set  $A_z = V_z = 0$ , use transverse gauge for  $V_\mu$  and decompose  $A_\mu$  on longitudinal and transverse parts.:

$$\partial_\mu V_\mu = 0 \quad A_\mu = A_{\perp\mu} + \partial_\mu \phi$$

One can relate the pseudoscalar current  $\bar{q}\gamma_5 q$  with axial vector current  $\bar{q}\gamma_5\gamma_\mu q$  via:

$$\partial_\mu (\bar{q}\gamma_5\gamma_\mu q) = 2m (\bar{q}\gamma_5 q)$$

So we can write out the following table of correspondence

$$\begin{aligned} V_\mu &\leftrightarrow \bar{q}\gamma^\mu q = J_V \\ A_\mu &\leftrightarrow \bar{q}\gamma_5\gamma^\mu q = J_A \\ \frac{Q^2}{2m}\phi &\leftrightarrow \bar{q}\gamma_5 q = J_\pi \end{aligned}$$

With this table, we can calculate two-point functions of QCD via our 5D theory, using the AdS/CFT recipe. For example:

$$\langle J_V(q_1)J_V(q_2) \rangle = \frac{\delta}{\delta V_0(q_1)} \frac{\delta}{\delta V_0(q_2)} S(V_{classical})|_{V_0=0}.$$

$$V_0(q) = V_{classical}(q, z)|_{z=0}$$

In order to perform this calculation one needs to find classical solutions, which are solutions to equations of motion, obtained by variation of  $S(V_\mu, A_\mu, \partial_\mu\phi, A_z)$ :

$$\begin{aligned} \left[ \partial_z \left( \frac{1}{z} \partial_z V_\mu^a \right) + \frac{q^2}{z} V_\mu^a \right]_\perp &= 0 \\ \left[ \partial_z \left( \frac{1}{z} \partial_z A_\mu^a \right) + \frac{q^2}{z} A_\mu^a - \frac{R^2 g_5^2 \Lambda^2 v^2}{z^3} A_\mu^a \right]_\perp &= 0 \\ \partial_z \left( \frac{1}{z} \partial_z \phi^a \right) + \frac{R^2 g_5^2 \Lambda^2 v^2}{z^3} (\pi^a - \phi^a) &= 0 \\ -q^2 \partial_z \phi^a + \frac{R^2 g_5^2 \Lambda^2 v^2}{z^2} \partial_z \pi^a &= 0 \end{aligned}$$

One can see, that  $X$  do not interact with vector field in quadratic action, so the equation for  $v$  is exactly solvable. Opposite to others, which can be solved perturbatively in the limits of large or small momenta.

### 3 Parameter fixing

Let's calculate some two-point functions. We start with vector current, associated with the field  $V$  in the model. The solution for  $V$  is:

$$V(Q, z) = -V_0(Q) \frac{1}{I_0(Qz_m)} Qz [K_0(Qz_m)I_1(Qz) - I_0(Qz_m)K_1(Qz)]$$

We substitute it to the variation of metric with respect to boundary value  $V_0$ , which can be presented in the form

$$\delta S_V = - \int d^4x \frac{R}{g_5^2} \left[ \delta V_\mu \frac{\partial_z V_\mu}{z} \right]_{z=\epsilon}$$

and the result for current correlator is:

$$\langle J_{V_\mu}^a(q) J_{V_\nu}^b(q) \rangle = \delta^{ab} (q_\mu q_\nu - q^2 g_{\mu\nu}) \Pi_V(q^2)$$

where

$$\Pi_V(Q^2) = -\frac{R}{g_5^2} \frac{\left( K_0(Qz_m) - I_0(Qz_m) [\ln(Q\epsilon/2) + \gamma] \right)}{I_0(Qz_m)}$$

The poles of Euclidean correlator correspond to masses of bound states. Consequently the first pole corresponds to the  $\rho$ -meson mass (the meson, associated with vector current in QCD), so we can fix the value of  $z_m$ :

$$I_0(iM_\rho z_m) = 0 \quad \implies \quad z_m = \frac{2.4}{M_\rho} = \frac{1}{323} \text{Mev}^{-1}$$

The masses of higher states we will obtain automatically: each one will correspond to the zero of Bessel function. The problem here is that this spectrum do not demonstrate the Regge behavior. In order to solve it, one have to change form of IR boundary. If at  $z_m$  metric has a factor of  $\exp(-z^2)$  instead of a simple cut, the Regge behavior will take place. [15, 16]

In the large  $Q^2$  limit we have

$$\Pi_V(Q^2) = -\frac{R}{2g_5^2} \ln Q^2 \epsilon^2$$

This result has the same form as in QCD and can be compared with the QCD sum rules leading term [18]:

$$\Pi_V(Q^2) = -\frac{N_c}{24\pi^2} \ln Q^2 \epsilon^2$$

And this fixes the 5D coupling constant  $g_5$

$$\frac{g_5^2}{R} = \frac{12\pi^2}{N_c}$$

To compute correlator of pseudoscalar current  $J_\pi$  we find solutions for coupled  $\phi$  and  $\pi$  near the boundary

$$\begin{aligned} \phi(z) &= \phi_0(q) Qz K_1(Qz). \\ \pi(z) &= -\phi_0(q) \frac{Q^2}{g_5^2 R^2 \Lambda^2 m^2} Qz K_1(Qz). \end{aligned}$$

The variation of action with respect to  $\phi_0(q)$  gives

$$\delta S_\pi = \int d^4x \frac{R}{g_5^2} \left[ \delta \partial_\mu \phi \frac{\partial_z \partial_\mu \phi}{z} \right]_{z=\epsilon} - \Lambda^2 R^3 \left[ \delta \pi \frac{v^2}{z^3} \partial_z \pi \right]_{z=\epsilon}$$

And we get for correlator:

$$\langle J_\pi(q), J_\pi(q) \rangle = 2 \frac{R}{g_5^2} \frac{1}{g_5^2 R^2 \Lambda^2} Q^2 \ln(Q^2 \epsilon^2)$$

One can find, that this result also has the same form as in QCD and comparison with the sum rules leading term [18]

$$\langle J_\pi(q), J_\pi(q) \rangle_{QCD} = \frac{N_c}{16\pi^2} Q^2 \ln(Q^2 \epsilon^2)$$

gives the value of  $\Lambda$

$$\Lambda^2 = \frac{8}{3} \frac{1}{g_5^2 R^2} = \frac{2N_c}{9\pi^2} \frac{1}{R^3}$$

We can also calculate the value of chiral condensate to fix the value of  $\sigma$ . As was mentioned above, it should be proportional to  $\sigma$ , but we need to find coefficient. In QCD the chiral condensate is defined as:

$$\langle \bar{q}q \rangle = \frac{\delta \epsilon_{QCD}}{\delta m_q} \Big|_{m_q=0}$$

On the AdS side this corresponds to:

$$\langle \bar{q}q \rangle = \frac{\delta S(X_0)}{\delta m} \Big|_{m=0} = 3R^3 \Lambda^2 \sigma = \frac{2N_c}{3\pi^2} \sigma$$

We see, that  $\langle \bar{q}q \rangle$  is proportional to  $\sigma$  and, because chiral condensate is linear in  $N_c$ ,  $\sigma$  turn out to be  $O(N_c^0)$ . For  $N_c = 3$   $\sigma = (462 \text{Mev})^3$ .

### 3.1 Results

After we have fixed all parameters in the model, we can write out the action in terms of QCD values:

$$S = \frac{N_c}{12\pi^2} \int d^5x \left\{ -\frac{1}{4z}(F_A^2 + F_V^2) + \frac{4}{3z^3}v(z)^2(\partial\pi - A)^2 + \frac{4}{z^5}v(z)^2 \right\}$$

We can calculate the axial two-point function. It is a little more difficult, than in vector case, because the equation of motion can't be solved exactly. We solve it in the limit of large  $Q^2$  perturbatively and corrections of order  $\frac{1}{Q^n}$  in the classical solution give next to leading terms of OPE. We can write out the result with corrections of order  $\frac{\sigma^2}{Q^6}$  and  $\frac{m\sigma}{Q^4}$ :

$$\Pi_A(Q^2) = -\frac{N_c}{24\pi^2} \left[ \ln Q^2 + \frac{128}{15} \frac{\sigma^2}{Q^6} - \frac{64}{9} \frac{\sigma m}{Q^4} \right]$$

One can see, that coefficients here do not coincide with sum rules calculation. We tune this result by adding a perturbation in the metric [13]:

$$ds^2 = \omega(z)(-dz^2 + dx^\mu dx_\mu), \quad \omega(z) = \frac{R^2}{z^2} + A\sigma^2 z^4 + B\sigma m z^2$$

which will cause the additional correction to the classical solution to appear, and this will change coefficients as desired.

The interesting object is "left-right" correlator  $\Pi_{LR} = \Pi_A - \Pi_V$

$$\Pi_{LR} = -\frac{N_c}{9\pi^2} \left[ \frac{16}{5} \frac{\sigma^2}{Q^6} - \frac{8}{3} \frac{\sigma m_q}{Q^4} \right]$$

Note here, that it has not powers of R, namely it has the order  $\lambda'^0$ , because in AdS/CFT:

$$\frac{R^4}{4\pi\alpha'^2} = \lambda' = N_c g_{ym}^2$$

If we denote coefficients in this formula as  $f$  and  $\rho$ , we find that at  $\lambda' \rightarrow \infty$  our calculation predicts:

$$f(\lambda') \sim \rho(\lambda') \sim \lambda'^0$$

while at weak coupling regime(sum rules):

$$\rho(\lambda') \sim \lambda'^0 \quad f(\lambda') = -4\pi\alpha_s \sim \lambda'.$$

The different behavior of the results is an evidence, that two approaches, AdS/QCD and sum rules, are applicable at different regimes. AdS/QCD works at at strong coupling, when sum rules are the calculation at week coupling.

One more value, that we can find, using AdS/QCD is  $f_\pi$ . We use the relation

$$\Pi_A(Q)|_{Q \rightarrow 0} = \frac{f_\pi^2}{Q^2}$$

We can solve the equation of motion of A in the limit  $Q^2 \rightarrow 0$  to obtain the two-point function at small  $Q^2$ . We find:

$$\begin{aligned} f_\pi^2 &= -\frac{R}{g_5} \frac{\partial_z a(z)}{z} \Big|_{z=0, Q=0} = \\ &\approx \frac{R}{g_5} 2.16\sigma^{2/3} = \frac{N_c}{12\pi^2} 2.16 \left( \frac{3\pi^2 \langle \bar{q}q \rangle}{2 N_c} \right)^{2/3} \sim 40 Mev \end{aligned}$$

This value obviously do not coincide with expected 140 Mev. We see, that  $f_\pi$  is related to the parameters of classical solution of X. If one introduce the additional potential for X in the 5D bulk. This solution will change and consequently the value of  $f_\pi$  can be tuned.

## 4 Conclusion

The model under consideration has several free parameters, but still has some predictive power. It gives qualitatively satisfactory results, but numbers differ. Study of such simple model gives an insight on common features of AdS/QCD and proposes modifications, needed to obtain more realistic results.

## 5 Acknowledgments

I am grateful to Alexander Gorsky for suggesting this problem and general guidance. I also would like to thank Konstantin Zarembo for useful discussions and Thomas Schaefer for important remarks.

I would like to adress the special thanks to the organizing committee of QUARKS-2008 for hospitality and interesting program of the seminar.

## References

- [1] A.Krikun, Phys.Rev.D77 No.12 (2008) [arXiv:(hep-th) 0801.4215]
- [2] J.M. Maldacena, Adv.Theor.Math.Phys. 2 (1998) 231-252 [arXiv:hep-th/9711200]; S.S. Gubser, I.R. Klebanov and A.M. Polyakov, Phys.Lett.B428, 105 (1998) [arXiv:hep-th/9802109]; E. Witten, Adv.Theor.Math.Phys. 2, 253 (1998). [arXiv:hep-th/9802150]
- [3] J. Erlich, E. Katz, D. T. Son, M. Stephanov, Phys.Rev.Lett. 95 (2005) 261602. [arXiv:hep-ph/0501128]
- [4] A. Karch, E.Katz, D.T. Son, M.Stephanov, Phys.Rev. D74 (2006) 015005. [arXiv:hep-ph/0602229]
- [5] L. Da Rold, A. Pomarol, Nucl.Phys. B721 (2005) 79-97. [arXiv:hep-ph/0501218]
- [6] L. Da Rold, A. Pomarol, JHEP 0601 (2006) 157. [arXiv:hep-ph/0510268]
- [7] N. Evans, A. Tedder, T.Watson, JHEP 0701 (2007) 058. [arXiv:hep-ph/0603249]
- [8] J. Erlich, G.D. Kribs, I. Low, Phys.Rev. D73 (2006) 096001. [arXiv:hep-th/0602110]
- [9] J. Polchinski, M.J. Strassler Phys.Rev.Lett.88:031601,2002. [arXiv: hep-th/0109174]
- [10] T.Sakai, S.Sugimoto, Prog.Theor.Phys. 113 (2005) 843-882 [arXiv:hep-th/0412141]
- [11] T.Sakai, S.Sugimoto, Prog.Theor.Phys. 114 (2006) 1083-1118. [arXiv: hep-th/0507073]
- [12] L. Randall and M. D. Schwartz, JHEP 0111, 003 (2001). [arXiv:hep-th/0108114].
- [13] J. Hirn, N. Rius, V. Sanz, Phys.Rev. D73 (2006) 085005. [arXiv:hep-ph/0512240]
- [14] S.J. Brodsky , G.F. de Teramond , Phys.Lett.B582:211-221,2004. [arXiv:hep-th/0310227]
- [15] A. Karch, E. Katz, D. T. Son and M. A. Stephanov, Phys. Rev. D 74, 015005 (2006). [arXiv:hep-ph/0602229]
- [16] O. Andreev, V.I. Zakharov, Phys. Rev. D 74, 025023 (2006). [arXiv:hep-ph/0604204]
- [17] J.M. Maldacena, H. Ooguri, Y. Oz et al., Phys.Rept.323:183-386,2000. [arXiv: hep-th/9905111]; I.R. Klebanov, Int.J.Mod.Phys.A21:1831-1843,2006. [arXiv: hep-ph/0509087]; A. Gorsky, Phys.Usp.48:1093-1108,2005. [arXiv: hep-th/0602184]

- [18] M.A. Shifman, A.I. Vainshtein, and V.I. Zakharov, Nucl. Phys. B147, 385, 448 (1979); V. A. Novikov, M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, Nucl. Phys. B191, 301 (1981)