

Precision of the Drell-Yan process description at LHC

Andrej Arbuzov

JINR, Dubna

Quarks 2010, 6-12 June, 2010

- Why Drell-Yan at LHC?
- SANC project
- Theoretical description of DY
- Electroweak corrections
- QCD corrections
- Theoretical uncertainties
- Conclusions

Motivation

- LHC is *not only a discovery* machine
- SM single W and Z production at LHC have *large cross sections* and *clear signatures*
- The experimental precision tag is **1%**
- CC and NC Drell-Yan processes will be used for
 - luminosity monitoring;
 - W mass and width measurement;
 - extraction of parton density functions;
 - detector calibration;
 - background to many other processes;
 - new physics searches
- **Are we ready to provide an adequately accurate theoretical description of Drell-Yan?**

SANC: a project to Support of Analytic and Numeric calculations for experiments at Colliders

- SANC web-site: <http://sanc.jinr.ru>, <http://pcphsanc.cern.ch>
- A. Andonov et al., “Standard SANC Modules,” Comput. Phys. Comm. **181** (2010) 305
- D. Bardin et al., “Implementation of SANC EW corrections in WINHAC Monte Carlo generator,” Acta Phys. Pol. B **40** (2009) 75
- A. Andonov et al., “NLO QCD corrections to Drell-Yan processes in the SANC framework,” Yad. Fiz. **73** (2010)
- D. Bardin et al. “J(A) functions in the Passarino-Veltman reduction,” Phys. Part. Nucl. Lett. **6** (2009) 30



Description of Drell-Yan like processes

Aiming at high precision of DY description we need:

- QCD in LO, NLO and NNLO
- Parton shower and hadronisation effects
- EW radiative corrections in one-loop at least
- Most important higher order effects (resummed where possible)
- Interplay of QCD and EW effects
- Input: coupling constants, hadronic vacuum polarization, and PDFs for the appropriate energy scales and x -values
- All relevant effects to be implemented in a Monte Carlo event generator(s)

Drell-Yan in SANC


Automatized analytic calculations in SANC provide FORM and FORTRAN modules, which can be downloaded and used in a stand-alone mode (see Comput. Phys. Comm.'09 and the SANC web-site)

For the CC and NC Drell-Yan in SANC we have:

- Complete one-loop EW RC [A.A. et al., EPJC'06 (CC DY); EPJC'08 (NC DY)]
- Photon induced DY processes [A.A. et al., JETP'08]
- Complete one-loop (NLO) QCD [A.Andonov et al., Yad. Fiz.'10; Phys.Part.Nucl.Lett.'07]
- Interface to parton showers in PYTHIA and HERWIG [P.Richardson, R.Sadykov, A.Sapronov, M.Seymour, P.Skands]
- Higher order photonic FSR in LLA
- MC integrator and MC event generator

Tuned comparison of EW RC

Tuned comparison with results of HORACE and Z(W)GRADE for EW RC to CC and NC DY were performed within *Les Houches* '05, '07 and *TEV4LHC* '06 workshops





- Set-up: $P_T(l, \nu) > 20$ GeV, $|\eta(l)| < 2.5$; $\alpha(0)$ EW scheme; MRST2004QED; NLO QED DIS subtraction scheme

Higher order photonic FSR

Multiple FSR photon radiation in LLA
Comparison with WINHAC (BARE e^+)

Preliminary




Interfacing SANC with PYTHIA and HERWIG

- SANC Monte-Carlo generators for neutral and charged current Drell-Yan processes produce unweighted events with help of FOAM algorithm
- The transfer of information between SANC Monte Carlo generator and the general purpose event generators PYTHIA and HERWIG is organized via data files containing the event information in the standard Les Houches Accord format
- Matching for the Born-level and $\mathcal{O}(\alpha)$ EW contributions with parton showers is *almost* trivial
- Matching of $\mathcal{O}(\alpha_s)$ QCD contributions is *scheme dependent*


The effect of parton showers on single-W production

Distribution of transverse momentum of μ^+




Distribution of Z transverse momentum

Distribution of transverse momentum of Z




Distribution of transverse momentum of Z




PS: PYTHIA vs HERWIG

SANC NC LO DY showered: Herwig++ vs Pythia8, $\alpha(0)$ -scheme, QED FSR off




SANC NC LO DY showered: Herwig++ vs Pythia8, $\alpha(0)$ -scheme, QED FSR off



QCD in NLO



- One-loop QCD corrections to Drell-Yan are computed in SANC for CC and NC case, including gluon induced sub-processes
- Comparison with MCFM is performed for CC and NC cases
- Matching of one-loop QCD RC in SANC MC with parton showers in PYTHIA and HERWIG is in progress. POWHEG scheme of matching is applied



Massive quark factorisation scheme (ACOT) in SANC

| | Born | NLO | δ |
|------------------|-------------|-----------|----------|
| CC [SANC], G_F | 2058.149(3) | 1906.1(8) | -7.39(4) |
| CC [MCFM], G_F | 2058.1(1) | 1901.9(9) | -7.59(5) |
| NC [SANC], G_F | 233.548(1) | 254.7(2) | 9.1(1) |
| NC [MCFM], G_F | 234.14(4) | 254.8(1) | 8.8(1) |

Choice of LO PDFs: NLO (CTEQ6M) vs LO



Drell-Yan: CC (left) and NC (right), $E_{CM} = 10$ TeV

Theoretical uncertainty

- Uncertainties in PDF, but after a new fit from LHC data ...
- Treatment of heavy quarks (c and b) ...
- QCD (and QED) factorization scheme and scale dependence
- pure QCD higher order terms \Leftarrow recent NNLO results by Anastasiou, Melnikov, Petriello *et al.* (FEWZ code)
- pure EW higher order terms:
 - EW scheme dependence: $\alpha(0)$ vs. G_F vs. $\alpha(M_Z)$
 - resummation of higher order EW Sudakov logs
 - other unknown EW higher order terms (should be small?)
 - hadronic vacuum polarization
 - top and Higgs mass dependence
- Interplay of EW and QCD effects: multiplicative vs. additive treatment
- All this to be quantified within working groups

- SANC provides an advanced description of CC and NC Drell-Yan processes
- Monte Carlo event generators are created, their development is continued
- Packages with MC and partonic level modules are available for download
- SANC modules with EW RC to CC Drell-Yan were implemented in WINHAC (ZINHAC soon) event generator
- Tuned comparison with other groups was performed, it should be extended
- But further theoretical studies are still required for better understanding of DY at LHC