



# Prompt photon production at HERA with $kt$ -factorization

S.P. Baranov<sup>1</sup>, A.V. Lipatov<sup>2</sup>, N.P. Zotov<sup>2</sup>

1 — Lebedev Physical Institute, Moscow, Russia

2 — SINP MSU, Moscow, Russia



# Content



---

- Motivation
- Phenomenology of  $kt$ -factorization
- Some calculation details
- Numerical results
  - photoproduction regime
  - DIS
- Summary

## Motivation I

---

- Prompt photons are directly coupled to the interacted quarks
- At HERA, they can be produced via direct and resolved photon events
- They directly probe the proton and photon PDFs
- They not affected by the subsequent fragmentation and hadronization



## Motivation II



- The NLO QCD predictions are typically 30 – 40% below the photoproduction data

A. Zembrzuski, M. Krawczyk,  
PRD 64, 114017 (2001)

M. Fontannaz, J.Ph. Guillet, G. Heinrich,  
EPJ C 21, 303 (2001)

- At DIS, there is substantial underestimation of the data at low  $Q^2$

A. Gehrmann-De Rider, G. Kramer,  
H. Spiesberger, PRL 96, 132006 (2006)







## Motivation III

---


- The visible disagreement in NLO calculations can be reduced by the introducing «by hands» of some intrinsic partonic transverse momentum  $k_t \sim 2 \text{ GeV}$ . However, such large  $k_t$  should have a significant QCD component
- The non-zero partonic  $k_t$  is naturally occurs in the ***kt-factorization approach of QCD***, where it is controlled by the non-collinear (BFKL-like) evolution equations
- First applications of the  $k_t$ -factorization approach to the prompt photon photo-production have been made

A.V. Lipatov, N.P. Zotov, PRD 72, 054002 (2005)

# Motivation IV



# kt-factorization approach I





## kt-factorization approach II

---

- Matrix elements of partonic subprocesses should be off-shell
- Partonic PDFs should be unintegrated (i.e. kt-dependent)
- Any observable can be calculated by the convolution of off-shell matrix elements with the unintegrated PDFs in both  $x$  and  $kt$


### See more details in

Small- $x$  Collaboration, EPJ C 48, 53 (2006); EPJ C 35, 67 (2004);  
EPJ C 25, 77 (2002)




# Calculations I

## Photoproduction



## DIS





## Calculations II

---

In off-shell matrix elements:

- Polarization tensor for off-shell gluons:  $\sum e^\mu e^\nu = \frac{k_T^\mu k_T^\nu}{k_T^2}$
- To calculate the spin density matrix for off-shell quarks, we extend the original diagram and consider the off-shell quark line as internal line in the extended diagram.
- In the small- $x$  approximation:  $\sum u \bar{u} = x \hat{p}_p$   
(neglecting also the quark masses)



## Calculations III

---

- **KMR unintegrated PDFs**

M.A. Kimber, A.D. Martin, M.G. Ryskin, PRD 63, 114027 (2001)

$$f_q(x, k_T^2, \mu^2) \sim T_q(k_T^2, \mu^2) [P_{qq} \times q(x, k_T^2) + P_{qg} \times g(x, k_T^2)]$$

$$f_g(x, k_T^2, \mu^2) \sim T_g(k_T^2, \mu^2) [P_{gq} \times q(x, k_T^2) + P_{gg} \times g(x, k_T^2)]$$

- **CCFM unintegrated PDFs**

gluon density has been fitted on DIS data H. Jung, arXiv:hep-ph/0411287

valence quark density has been proposed M. Deak, H. Jung, K. Kutak, DIS'08

sea quark density is approximated by the last gluon splitting, i.e.

$$f_q^{(s)}(x, k_T^2, \mu^2) \sim P_{qg} \times f_g(x, k_T^2, \mu^2)$$



## Calculations IV

---



Numerical parameters:

- Massless approximation for all quark flavours
- Isolation criterion (also removes the fragmentation photons)
- $N_f = 4$ ,  $\Lambda_{QCD} = 200 \text{ MeV}$
- Hard scale  $\mu^2 = E_T^2$

# Numerical results: incl. photoproduction

Solid histograms — KMR uPDFs  
 Dashed (dash-dotted) — CCFM set A0 (B0)



Missing sea quarks component  
 in CCFM



# Numerical results: incl. photoproduction

Solid histograms — KMR uPDFs  
 Dashed (dash-dotted) — CCFM set A0 (B0)

Missing sea quarks component  
 in CCFM







# Numerical results: excl. photoproduction

Soild histograms — KMR uPDFs

Dashed (dash-dotted) — CCFM set A0 (B0)







# Numerical results: excl. photoproduction

$$p_T = E_T^\gamma \sin \Delta \varphi$$

Sensitive to the high-order contributions

Solid histograms — KMR

Dashed (dash-dotted) — CCFM







# Numerical results: DIS

Soild histograms — KMR uPDFs

Upper and lower dashed histograms — scale variations in KMR







# Numerical results: DIS

Soild histograms — KMR uPDFs



Upper and lower dashed histograms — scale variations in KMR



# Numerical results: DIS



Soild histograms — KMR uPDFs

Upper and lower dashed histograms — scale variations in KMR



# Numerical results: DIS

Different description of the H1 and ZEUS data  
in the exclusive production case





## Summary

---

- We presented the numerical calculations for the prompt photon production at HERA (both photo-production and DIS)
- We find a good agreement with the HERA data
- The transverse momentum of initial partons is important for description of the HERA data
- The higher-order QCD contributions are effectively simulated in the  $kt$ -factorization approach at LO level
- The contribution from the quarks are important and should be included into the non-collinear evolution cascade