# Four problems for the $c - \tau$ , b and super- $c - \tau$ , b factories

N.N. Achasov, A.V. Kiselev, A.A. Kozhevnikov, E.V. Rogozina, and G.N. Shestakov

Laboratory of Theoretical Physics,

**Sobolev Institute for Mathematics,** 

Academician Koptiug Prospekt, 4, Novosibirsk, 630090, Russia

Electronic Address: achasov@math.nsc.ru

QUARKS-2014, June 2-8, 2014, Suzdal – p.1/51

### ABSTACT

The problems suggested by authors for studying at the  $c-\tau$ , b and super- $c-\tau$ , b factories are stated.

• Comparison the production mechanism of the light scalar mesons in  $D_s^+ \to s\bar{s} e^+ \nu \to [\sigma(600) + f_0(980)] e^+ \nu \to \pi^+ \pi^- e^+ \nu$ with the production mechanism of the light pseudoscalar mesons in  $D_s^+ \to s\bar{s} e^+ \nu \to (\eta/\eta') e^+ \nu$  shows that  $s\bar{s} \to \sigma(600)$  is negligibly small in comparison with  $s\bar{s} \to f_0(980)$ . As for  $f_0(980), s\bar{s} \to f_0(980)$  is not more 30% of  $s\bar{s} \to \eta_s$  ( $\eta_s = s\bar{s}$ ).

The study of the light scalar mesons in semileptonic decays of the  $D^+(D^-), D^0(\bar{D}^0), B^+(B^-), B^0(\bar{B}^0)$  mesons is suggested.

QUARKS-2014, June 2-8, 2014, Suzdal – p.2/51

# ABSTACT

• Interference phenomenon observed in the  $\psi(3770)$  resonance region the  $e^+e^- \rightarrow D\bar{D}$  reactions is described with models satisfying the elastic unitarity requirement. As a candidate, a model with the mixing  $\psi(3770)$  and  $\psi(2S)$  resonances is proposed. The selection of theoretical models in the non- $D\bar{D}$  channels  $e + e - \rightarrow \psi(3770) \rightarrow \gamma \chi_{c0}, J/\psi\eta, \phi\eta$ , etc is suggested.

• • The branching ratios (BR) of decays  $\psi(3770)$  and  $\Upsilon(10580)$ into light (non- $D\bar{D}$  and non- $B\bar{B}$ ) hadrons caused by the intermediate real  $D\bar{D}$  and  $B\bar{B}$  states are calculated. We got a band of predictions for the branching ratios : 1%  $\leq$  BR  $\leq$  15%. The lower bound is 10 times as large as the branching ratio of annihilation into three gluons.

QUARKS-2014, June 2-8, 2014, Suzdal – p.3/51

### ABSTACT

• • • Enfant terrible of charmonium spectroscopy, the resonance X(3872), generated a stream of interpretations and ushered in a new exotic XYZ spectroscopy. In the meantime, many (if not all) characteristics of X(3872) are rather ambiguous. We construct spectra of decays of X(3872) with good analytical and unitary properties which allows to define the branching ratio of the  $X(3872) \rightarrow D^{*0}\bar{D}^0 + c.c.$  decay studying only one more decay, for example, the  $X(3872) 
ightarrow \pi^+\pi^- J/\psi(1S)$  decay. We next define the range of values of the coupling constant of X(7872) with the  $D^{*0}\overline{D}^0$  system. Finally, we show that our spectra are effective means of selection of models for X(3872).

QUARKS-2014, June 2-8, 2014, Suzdal – p.4/51

Light scalars in semi-leptonic decays of heavy quarkonia

Based on N.N.Achasov and A.V. Kiselev, Physical Review D 86, 114010 (2012)

It is time to explore the light scalar mesons in the decays of heavy quarkonia. The semi-leptonic decays are of prime interest because they have the clear mechanisms.

QUARKS-2014, June 2-8, 2014, Suzdal – p.5/51



Model of the  $D^+_s o (\sigma/f_0) \, e^+ 
u$  and  $D^+_s o (\eta/\eta') \, e^+ 
u$  decays

QUARKS-2014, June 2-8, 2014, Suzdal – p.6/51

### The $D_s^+ o (\sigma/f_0) \, e^+ u$ and $D_s^+ o (\eta/\eta') \, e^+ u$ decays

Below we study the mechanism of production of the light scalar mesons in the  $D_s^+ 
ightarrow \pi^+\pi^- \, e^+ 
u$  decays:  $D_s^+ \to s\bar{s} e^+ \nu \to [\sigma(600) + f_0(980)] e^+ \nu \to \pi^+ \pi^- e^+ \nu,$ and compare it with the mechanism of production of the light pseudoscalar mesons in the  $D_s^+ 
ightarrow (\eta/\eta') \, e^+ 
u$  decays:  $D_s^+ o s \overline{s} \, e^+ 
u o (\eta/\eta') \, e^+ 
u$ , in a model of the NJL type.  $M[D_s^+(p) 
ightarrow P(p_1)W^+(q) 
ightarrow P(p_1)e^+
u] = rac{G_F}{\sqrt{2}}V_{cs}V_{lpha}L^{lpha}$  $M[D_s^+(p)
ightarrow S(p_1)W^+(q)
ightarrow S(p_1)e^+
u]=rac{G_F}{\sqrt{2}}V_{cs}A_lpha L^lpha\,,$  $V_{\alpha} = f^{P}_{+}(q^{2})(p + p_{1})_{\alpha} + f^{P}_{-}(q^{2})(p - p_{1})_{\alpha},$  $A_{\alpha} = f_{+}^{S}(q^{2})(p + p_{1})_{\alpha} + f_{-}^{S}(q^{2})(p - p_{1})_{\alpha},$  $L_{\alpha} = \bar{\nu} \gamma_{\alpha} (1 + \gamma_5) e$ ,  $q = (p - p_1)$ .

The influence of  $f_-^P(q^2)$  and  $f_-^S(q^2)$  are negligible for  $m_{e^+}.$ 

QUARKS-2014, June 2-8, 2014, Suzdal – p.7/51

#### The decay rates in the stable *P* and *S* states

$$rac{d\Gamma(D_s^+ o P\,e^+
u)}{dq^2} = rac{G_F^2 |V_{cs}|^2}{24\pi^3} p_1^3(q^2) |f_+^P(q^2)|^2, 
onumber \ rac{d\Gamma(D_s^+ o S\,e^+
u)}{dq^2} = rac{G_F^2 |V_{cs}|^2}{24\pi^3} p_1^3(q^2) |f_+^S(q^2)|^2.$$

For the  $f^P_+(q^2)$  and  $f^S_+(q^2)$  form factors we use the vector dominance model

$$egin{aligned} f^P_+(q^2) &= f^P_+(0) rac{m_V^2}{m_V^2 - q^2} = f^P_+(0) f_V(q^2)\,, \ f^S_+(q^2) &= f^S_+(0) rac{m_A^2}{m_A^2 - q^2} = f^S_+(0) f_A(q^2)\,, \end{aligned}$$

where  $V = D_s^*(2112)^\pm$  ,  $A = D_{s1}(2460)^\pm$  .

QUARKS-2014, June 2-8, 2014, Suzdal – p.8/51

### **Definitions**

Following the NJL type model we write  $f^P_+(0)$  and  $f^S_+(0)$  in the form

$$f^P_+(0) = g_{D^+_s c ar s} F_P g_{s ar s P} \,, \quad f^S_+(0) = g_{D^+_s c ar s} F_S g_{s ar s S} \,.$$

We know the structure of  $\eta$  and  $\eta'$ 

 $\eta = \eta_q \cos \phi - \eta_s \sin \phi$ ,  $\eta' = \eta_q \sin \phi + \eta_s \cos \phi$ ,

where  $\eta_q = (u ar{u} + d ar{d})/\sqrt{2}$  and  $\eta_s = s ar{s}$ .

The angle  $\phi = \theta_i + \theta_P$ , where  $\theta_i$  is the ideal mixing angle with  $\cos \theta_i = \sqrt{1/3}$  and  $\sin \theta_i = \sqrt{2/3}$ , i.e.,  $\theta_i = 54.7^\circ$ , and  $\theta_P$  is the angle between the flavor-singlet state  $\eta_1$  and the flavor-octet state  $\eta_8$ .

QUARKS-2014, June 2-8, 2014, Suzdal – p.9/51

### **Definitions**

Particle Data Group give the  $\theta_P$  band  $-20^\circ \lesssim \theta_P \lesssim -10^\circ$  that gives us the opportunity to extract information about the  $s\bar{s} \to \eta_s$ coupling constant,  $g_{s\bar{s}}\eta_s$ , from experiment and to compare with the  $s\bar{s} \to f_0$  coupling constant,  $g_{s\bar{s}}f_0$ , extracted from experiment also.

#### We consider the next set of $\theta_P$ .

$$egin{aligned} & heta_P = -11^\circ : & \eta = 0.72\eta_0 - 0.69\eta_s \,, & \eta' = 0.69\eta_0 + 0.72\eta_s \ & heta_P = -14^\circ : & \eta = 0.76\eta_0 - 0.65\eta_s \,, & \eta' = 0.65\eta_0 + 0.76\eta_s \ & heta_P = -18^\circ : & \eta = 0.8\eta_0 - 0.6\eta_s \,, & \eta' = 0.6\eta_0 + 0.8\eta_s \,. \end{aligned}$$

$$BR(D_s^+ \to s\bar{s} e^+ \nu \to \eta e^+ \nu) = (2.67 \pm 0.29)\%,$$
  
 $BR(D_s^+ \to s\bar{s} e^+ \nu \to \eta' e^+ \nu) = (9.9 \pm 2.3) \times 10^{-3}.$ 

QUARKS-2014, June 2-8, 2014, Suzdal – p.10/51

$$\begin{split} D_{s}^{+} &\rightarrow s \overline{s} \ e^{+} \nu \ \rightarrow \pi^{+} \pi^{-} \ e^{+} \nu \\ \hline M(D_{s}^{+} \rightarrow s \overline{s} \ e^{+} \nu \rightarrow [\sigma(600) + f_{0}(980)] \ e^{+} \nu \rightarrow \pi^{+} \pi^{-} \ e^{+} \nu) \\ &= \frac{G_{F}}{\sqrt{2}} V_{cs} \ L^{\alpha} \ (p + p_{1})_{\alpha} \ g_{D_{s}^{+} c \overline{s}} \ f_{A}(q^{2}) \times \\ e^{i \delta_{B}^{\pi \pi}} \frac{1}{\Delta(m)} \left( F_{\sigma} g_{s \overline{s} \sigma} D_{f_{0}}(m) g_{\sigma \pi^{+} \pi^{-}} + F_{\sigma} g_{s \overline{s} \sigma} \Pi_{\sigma f_{0}}(m) g_{f_{0} \pi^{+} \pi^{-}} \right) \\ &+ F_{f_{0}} g_{s \overline{s} f_{0}} \Pi_{f_{0} \sigma}(m) g_{\sigma \pi^{+} \pi^{-}} + F_{f_{0}} g_{s \overline{s} f_{0}} D_{\sigma}(m) g_{f_{0} \pi^{+} \pi^{-}} \right), \\ &\text{where } m \text{ is the invariant mass of the } \pi \pi \text{ system, } \Delta(m) = \\ D_{f_{0}}(m) D_{\sigma}(m) - \Pi_{f_{0} \sigma}(m) \Pi_{\sigma f_{0}}(m), \ D_{\sigma}(m) \text{ and } D_{f_{0}}(m) \\ &\text{are the inverted propagators of the } \sigma \text{ and } f_{0} \text{ mesons, } \Pi_{\sigma f_{0}}(m) = \\ \Pi_{f_{0} \sigma}(m) \text{ is the off-diagonal element of the polarization operator,} \\ &\text{which mixes the } \sigma \text{ and } f_{0} \text{ mesons.} \end{split}$$

QUARKS-2014, June 2-8, 2014, Suzdal – p.11/51

 $D^+_s o \pi^+\pi^- \, e^+
u$ 

$$\begin{aligned} \frac{d^2 \Gamma(D_s^+ \to \pi^+ \pi^- e^+ \nu)}{dq^2 dm} &= \frac{G_F^2 |V_{cs}|^2}{24 \pi^3} g_{D_s^+ c\bar{s}}^2 |f_A(q^2)|^2 p_1^3(q^2, m) \\ &\times \frac{1}{8\pi^2} m \rho_{\pi\pi}(m) \left|\frac{1}{\Delta(m)}\right|^2 \end{aligned}$$

$$imes \Big| F_\sigma g_{sar{s}\sigma} D_{f_0}(m) g_{\sigma\pi^+\pi^-} + F_\sigma g_{sar{s}\sigma} \Pi_{\sigma f_0}(m) g_{f_0\pi^+\pi^-}$$

$$+ \, F_{f_0} g_{s ar{s} f_0} \Pi_{f_0 \sigma}(m) g_{\sigma \pi^+ \pi^-} + F_{f_0} g_{s ar{s} f_0} D_\sigma(m) g_{f_0 \pi^+ \pi^-} \Big|^2 \, ,$$

where 
$$ho_{\pi\pi}(m)=\sqrt{1-4m_\pi^2/m^2}$$
 .

QUARKS-2014, June 2-8, 2014, Suzdal – p.12/51



QUARKS-2014, June 2-8, 2014, Suzdal – p.13/51

# The $q^2$ distribution, the CLEO data



The  $q^2$  distribution for  $BR(D_s^+ \to f_0(980) e^+ \nu)$ . The axialvector dominance model (the theoretical curve) describes the data quite satisfactorily.

QUARKS-2014, June 2-8, 2014, Suzdal – p.14/51

# **Results of the analysis of the CLEO data**

$Br(D_s^+  ightarrow f_0 e^+  u  ightarrow \pi^+ \pi^- e^+  u) = 0.17\%$									
$\frac{F_{\sigma}g_{s\bar{s}\sigma}}{F_{f_0}g_{s\bar{s}f_0}}$	$\frac{F_{f_0}^2g_{s\bar{s}f_0}^2}{F_{\eta}^2g_{s\bar{s}\eta}^2}$	$\frac{F_{f_0}^2 g_{s\bar{s}f_0}^2}{F_{\eta'}^2 g_{s\bar{s}\eta'}^2}$	$rac{F_{\eta}^2 g_{sar{s}\eta}^2}{F_{\eta^\prime}^2 g_{sar{s}\eta^\prime}^2}$						
0.039	0.67	0.49	0.73						
The $\eta-\eta'$ mixing									
$ heta_P$	$-11^{\circ}$	$-14^{\circ}$	$-18^{\circ}$						
$rac{F_{f_0}^2g_{sar{s}f_0}^2}{F_\eta^2g_{sar{s}\eta_s}^2}$	0.32	0.29	0.24						
$\frac{F_{f_0}^2 g_{s\bar{s}f_0}^2}{F_{\eta'}^2 g_{s\bar{s}\eta_s}^2}$	0.27	0.28	0.31						

QUARKS-2014, June 2-8, 2014, Suzdal – p.15/51

### **Discussion and conclusion**

When fitting the CLEO data, we use the parameters of the resonances obtained by us in PRD 85, 094016 (2012) in the analysis of the  $\pi\pi$  scattering and the  $\phi \rightarrow \gamma(\sigma + f_0) \rightarrow \gamma\pi^0\pi^0$  decay. In addition, we take into account the Adler self consistency condition (the Adler zero at  $m^2$  near  $(m_{\pi}^2)/2$ ). Fitting the shape we fix only one parameter  $f_{+}^{\sigma}(0)/f_{+}^{f_0}(0) = F_{\sigma}g_{s\bar{s}\sigma}/F_{f_0}g_{s\bar{s}f_0}$ =0.039, 0.014, 0.055, 0.058, 0.032, 0.055 for six fits from PRD 85, 094016 (2012).

The 44 events in Fig. on page 13 determine only one parameter  $f^{\sigma}_{+}(0)/f^{f_0}_{+}(0)$ . The branching ratio fixes  $f^{f_0}_{+}(0)$ .

So the intensity of the  $\sigma(600)$  production is much less than the intensity of the  $f_0(980)$  production ( $(f_+^{\sigma}(0)/f_+^{f_0}(0))^2 < 0.003$ ).

QUARKS-2014, June 2-8, 2014, Suzdal – p.16/51

### **Discussion and conclusion**

That is we find the direct evidence of decoupling of  $\sigma(600)$  with the  $s\bar{s}$  pair. As far as we know, this is truly a new result, which agrees well with the decoupling of  $\sigma(600)$  with the  $K\bar{K}$  states, obtained in PRD 85, 094016 (2012)

 $g^2_{\sigma K^+K^-}/g^2_{\sigma \pi^+\pi^-}$  =0.04, 0.001, 0.01, 0.01, 0.003, 0.025 for six fits.

The decoupling of  $\sigma(600)$  with the  $K\bar{K}$  states means also the decoupling of  $\sigma(600)$  with  $\sigma_q = (u\bar{u} + d\bar{d})/\sqrt{2}$  because  $\sigma_q$  results in  $g^2_{\sigma K^+K^-}/g^2_{\sigma \pi^+\pi^-} = 1/4$ .

Fit 1 describes the  $\pi^+\pi^-$  spectrum better than others,  $(f^{\sigma}_+(0)/f^{f_0}_+(0))^2 = (0.039)^2$ ,  $g^2_{\sigma K^+K^-}/g^2_{\sigma \pi^+\pi^-} = 0.04$ .

So,the CLEO experiment gives new support in favour of the fourquark,  $ud\bar{u}d\bar{d}$ , structure of the  $\sigma(600)$  meson.

QUARKS-2014, June 2-8, 2014, Suzdal – p.17/51

### **Discussion and conclusion**

In the chirally symmetric model of the NJL type the coupling constants of the pseudoscalar and scalar partners with quarks are equal to each other, i.e.,  $g_{s\bar{s}\eta_s} = g_{s\bar{s}f_{0s}}$ , where  $f_{0s} = s\bar{s}$ . If to neglect the strange quark mass as compared with the charmed quark mass ( $m_s/m_c \ll 1$ ) in the numerators of the integrands for the decay diagrams, then  $F_{f_0} = F_{\eta'}$  and we find that  $g_{s\bar{s}f_0}^2/g_{s\bar{s}\eta_s}^2 \approx 0.3$ . So, the  $f_{0s} = s\bar{s}$  part in the  $f_0(980)$  wave function is near thirty percent.

Taking into account the suppression of the  $f_0(980)$  meson coupling with the  $\pi\pi$  system,  $g_{f_0\pi^+\pi^-}^2/g_{f_0K^+K^-}^2 = 0.154$ , one can conclude that the  $f_{0q} = (u\bar{u} + d\bar{d})/\sqrt{2}$  part in the  $f_0(980)$ wave function is suppressed also.

So, the CLEO experiment gives new support in favour of the fourquark,  $(sd\bar{s}d\bar{d} + sd\bar{s}d\bar{d})/\sqrt{2}$ , structure of the  $f_0(980)$  meson, too. QUARKS-2014, June 2-8, 2014, Suzdal – p.18/51

# Outlook

Certainly, there is an extreme need in experiment on the  $D_s^+ \to s\bar{s} \, e^+ \nu \to \pi^+ \pi^- \, e^+ \nu$  decay with high statistics.

Of great interest is the experimental search for the decays  $D^0 
ightarrow d\bar{u} e^+ \nu 
ightarrow a_0^-(980) e^+ \nu 
ightarrow \pi^- \eta e^+ \nu$  and  $D^+ 
ightarrow d\bar{d} e^+ \nu 
ightarrow a_0^0(980) e^+ \nu 
ightarrow \pi^0 \eta e^+ \nu$  (or the charge conjugate ones), which will give the information about the  $a_q^- = d\bar{u}$  (or  $a_q^+ = u\bar{d}$ ) component in the  $a_0^-(980)$ (or  $a_0^+(980)$ ) wave function and  $a_q^0 = (u\bar{u} - d\bar{d})/\sqrt{2}$ component in the  $a_0^0$  wave function.

#### Now it is known that

 $BR(D^0 \to d\bar{u} \, e^+ \nu \to \pi^- \, e^+ \nu) = (2.89 \pm 0.08) \times 10^{-3}$  and  $BR(D^+ \to d\bar{d} \, e^+ \nu \to \pi^0 \, e^+ \nu) = (4.05 \pm 0.18) \times 10^{-3}.$ 

QUARKS-2014, June 2-8, 2014, Suzdal – p.19/51

# Outlook

No less interesting is also search for the decays  $D^+ \rightarrow d\bar{d} \, e^+ \nu \rightarrow [\sigma(600) + f_0(980)] \, e^+ \nu \rightarrow \pi^+ \pi^- e^+ \nu$  (or the charge conjugate ones), which will give the information about the  $\sigma_q = (u\bar{u} + d\bar{d})/\sqrt{2}$  and  $f_{0q} = (u\bar{u} + d\bar{d})/\sqrt{2}$  components in the  $\sigma(600)$  and  $f_0(980)$  wave functions respectively.

Now it is known that  $BR(D^+ \to d\bar{d} e^+ \nu \to \eta e^+ \nu) = (1.14 \pm 0.10) \times 10^{-3}$  and  $BR(D^+ \to d\bar{d} e^+ \nu \to \eta' e^+ \nu) = (2.2 \pm 0.5) \times 10^{-4}$ .

Comparative research of light scalar and pseudoscalar mesons in semileptonic decays of B quarkonia at super B-factories is very tempting. Now it is known that

QUARKS-2014, June 2-8, 2014, Suzdal – p.20/51

### Outlook

$$\begin{split} BR(B^0 &\to d\bar{u} \, e^+\nu \to \pi^- \, e^+\nu) = (1.44 \pm 0.05) \times 10^{-4}, \\ BR(B^+ &\to u\bar{u} \, e^+\nu \to \pi^0 \, e^+\nu) = (7.79 \pm 0.26) \times 10^{-5}, \\ BR(B^+ &\to u\bar{u} \, e^+\nu \to \eta \, e^+\nu) = (3.8 \pm 0.6) \times 10^{-5} \text{ and} \\ BR(B^+ &\to u\bar{u} \, e^+\nu \to \eta' \, e^+\nu) = (2.3 \pm 0.8) \times 10^{-5}. \end{split}$$

#### QUARKS-2014, June 2-8, 2014, Suzdal – p.21/51

#### Interference phenomena in the $\psi(3770)$ resonance region

Based on N.N. Achasov and G.N. Shestakov, Physical Review D 86, 114013 (2012) and Physical Review D 87, 057502 (2013)



QUARKS-2014, June 2-8, 2014, Suzdal – p.22/51

#### The D meson electromagnetic form factor $F_D^0$

A similar representation of the  $e^+e^- \rightarrow D\bar{D}$  reaction amplitude used for the data description guarantees the unitarity requirement on the model level.

The sum of the  $e^+e^- \to D^0 \bar{D}^0$  and  $e^+e^- \to D^+D^-$  reaction cross sections is expressed in terms of  $F_D^0$  in the following way

$$\sigma(\gamma\gamma o Dar{D}) = rac{8\pilpha^2}{3s^2} \left|F_D^0(s)
ight|^2 \ 
u(s),$$

where  $\nu(s) = [p_0^3(s) + p_+^3(s)]/\sqrt{s}$ ,  $p_{0,+}(s) = \sqrt{s/4} - m_{D^{0,+}}^2$ .

QUARKS-2014, June 2-8, 2014, Suzdal – p.23/51

The model for  $F_D^0$  with the mixing  $\psi''$  and  $\psi(2S)$  resonances

It is clear that the main sources of the background in the  $\psi''$  region are the tails from the  $J/\psi$ ,  $\psi(2S)$ ,  $\psi(4040)$ ,  $\psi(4160)$  and other resonances. It is easy to incorporate the right number of resonances in our scheme.

Here we present the simplest variant of the model taking into account the background contribution from the nearest neighbor resonance  $\psi(2S)$  and also discuss how it can be checked.

In the considered model the  $\psi''$  and  $\psi(2S)$  resonances mix via transitions  $\psi'' o D\bar{D} o \psi(2S)$ .

QUARKS-2014, June 2-8, 2014, Suzdal – p.24/51

The model for  $F_D^0$  with the mixing  $\psi''$  and  $\psi(2S)$  resonances

$$F_D^0(s) = \frac{\mathcal{R}_{D\bar{D}}(s)}{D_{\psi''}(s)D_{\psi(2S)}(s) - \Pi^2_{\psi''\psi(2S)}(s)},$$

$$where \ \ D_{\psi^{\prime\prime}}(s) = m_{\psi^{\prime\prime}}^2 - s - i \sqrt{s} \Gamma_{\psi^{\prime\prime} D ar{D}}(s),$$

$$D_{\psi(2S)}(s) = m_{\psi(2S)}^2 - s - i\sqrt{s}\Gamma_{\psi(2S)D\bar{D}}(s),$$

$$\Gamma_{\psi''D\bar{D}}(s) = \frac{g_{\psi''D\bar{D}}^2}{6\pi} \frac{\nu(s)}{\sqrt{s}}, \ \Gamma_{\psi(2S)D\bar{D}}(s) = \frac{g_{\psi(2S)D\bar{D}}^2}{6\pi} \frac{\nu(s)}{\sqrt{s}},$$

the  $\psi'' - \psi(2S)$  mixing amplitude caused by  $\psi'' o D\bar{D} o \psi(2S)$  transitions via the real  $D\bar{D}$  intermediate states has the form

$$\Pi_{\psi''\psi(2S)}(s) = i g_{\psi''D\bar{D}} g_{\psi(2S)D\bar{D}} \nu(s)/(6\pi) ,$$

 $\mathcal{R}_{D\bar{D}}(s) = (m_{\psi''}^2 - s)g_{\psi(2S)\gamma}g_{\psi(2S)D\bar{D}} + (m_{\psi(2S)}^2 - s)g_{\psi''\gamma}g_{\psi''D\bar{D}}.$ 

#### QUARKS-2014, June 2-8, 2014, Suzdal – p.25/51

The model for  $F_D^0$  with the mixing  $\psi''$  and  $\psi(2S)$  resonances

 $m_{\psi''}, \ g_{\psi''D\bar{D}}, \ g_{\psi''\gamma}$ , and  $\ g_{\psi(2S)D\bar{D}}$  are determined by fitting;  $m_{\psi(2S)}$  and  $g_{\psi(2S)\gamma}$  are fixed by the PDG data.

Note that  $F_D^0$  in the considered model is proportional to the firstdegree polynomial in s with real coefficients (see  $\mathcal{R}_{D\bar{D}}(s)$  above ). Hence the dip observed in  $\sigma(e^+e^- \rightarrow D\bar{D})$  near 3.81 GeV can be explained by the  $F_D^0(s)$  zero, caused by compensation between the  $\psi''$  and  $\psi(2S)$  contributions.

QUARKS-2014, June 2-8, 2014, Suzdal – p.26/51

The simplest variant of the  $\psi^{\prime\prime}-\psi(2S)$  mixing model for  $F_D^0$ 



The solid curve is the fit to the data. The dashed and dot-dashed curves show the  $\psi''$  and  $\psi(2S)$  contributions, respectively. Bare parameters:  $m_{\psi''}$  = 3.794 GeV,  $\Gamma_{\psi'' D\bar{D}}$  = 56.8 MeV,  $\Gamma_{\psi'' e^+ e^-}$  = 0.062 keV,  $g^2_{\psi(2S)D\bar{D}}/4\pi$  = 32.2.

QUARKS-2014, June 2-8, 2014, Suzdal – p.27/51

The simplest variant of the  $\psi^{\prime\prime}-\psi(2S)$  mixing model for  $T_1^0$ 

From the fitting of the  $e^+e^- \rightarrow D\bar{D}$  data we all know, at the model level, about the I=0 P wave  $D\bar{D}$  elastic scattering amplitude  $T_1^0$ :

$$T_1^0(s) = e^{i\delta_1^0(s)} \sin \delta_1^0(s) = 
onumber \ = rac{
u(s)}{6\pi} \left[ rac{(m_{\psi''}^2 - s)g_{\psi(2S)Dar D}^2 + (m_{\psi(2S)}^2 - s)g_{\psi''Dar D}^2}{D_{\psi''}(s)D_{\psi(2S)}(s) - \Pi_{\psi''\psi(2S)}^2(s)} 
ight] \,.$$

QUARKS-2014, June 2-8, 2014, Suzdal – p.28/51

#### Cross section and phase for $Dar{D}$ elastic scattering in the P wave



(a) The cross section  $\sigma(D^0 \overline{D}^0 \to D^0 \overline{D}^0) = 3\pi |\sin \delta_1^0(s)|^2 / p_0^2(s)$  and (b) the phase  $\delta_1^0(s)$  for the simplest variant of the  $\psi'' - \psi(2S)$  mixing model. Unfortunately, these predictions are not possible to verify. However, there are many other reactions which can be measured experimentally.

QUARKS-2014, June 2-8, 2014, Suzdal – p.29/51

#### The $\psi''$ shapes in non- $Dar{D}$ decay channels



The solid curves show predictions of the model with the mixing  $\psi''$  and  $\psi(2S)$ resonances for the  $\psi''$  peak shapes in the  $e^+e^- \rightarrow \gamma \chi_{c0}$ ,  $e^+e^- \rightarrow J/\psi\eta$ , and  $e^+e^- \rightarrow \phi\eta$  cross sections; the dashed and dotted curves show the contributions from  $\psi''$  and  $\psi(2S)$  production amplitudes proportional to  $g_{\psi''ab}$  and  $g_{\psi(2S)ab}$ , respectively ( $ab = \gamma \chi_{c0}$ ,  $J/\psi\eta$ ,  $\phi\eta$ ). The points with errors are the CLEO data.

QUARKS-2014, June 2-8, 2014, Suzdal – p.30/51

#### Conclusion

- 1. The  $\psi''$  resonance shape keep important information about the production mechanism and interference with background. Its description requires taking into account the unitarity.
- 2. The simplest model mixing the  $\psi''$  and  $\psi(2S)$  resonances satisfies the unitarity requirement and describes the current data on the  $e^+e^- \rightarrow D\bar{D}$  reaction cross section very well. We also extracted from experiment  $g^2_{\psi(2S)D\bar{D}}/4\pi \approx 13 - 30$ .
- 3. New high-statistics data on the reactions  $e^+e^- \rightarrow D\bar{D}$  should help reveal the complex mechanism of the  $\psi''$  production.
- 4. The measurements of mass spectra in the  $\psi''$  region in the non- $D\bar{D}$  channels, such as  $e^+e^- \rightarrow \gamma \chi_{c0}$ ,  $J/\psi\eta$ ,  $\phi\eta$ , etc., will promote comprehensive study of the  $\psi''$  resonance physics and effective selection of theoretical models.

QUARKS-2014, June 2-8, 2014, Suzdal – p.31/51

Branching ratios of decays  $\psi(3770), \psi(4040), \text{ and } \Upsilon(10580)$  into light hadrons

Based on N.N. Achasov and A.A. Kozhevnikov, Phys. Rev. D 49, 275 (1994) and Yad. Fiz. 69, 1017 (2006) [Phys. At. Nucl. 69, 988 (2006)].

Exclusive decays of the ground-state  $c\bar{c}$  and bb quarkonia  $J/\psi(1S)$  and  $\Upsilon(1S)$  into light hadrons are qualitatively similar in that their branching ratios are very small,  $\sim 10^{-3} - 10^{-4}$ . Since in the framework of the quark-gluon picture such decays are originated from the 3-gluon annihilation, a rough estimate gives  $\Gamma((Q\bar{Q} \rightarrow (q\bar{q}) + (q\bar{q})) \sim \alpha_s^3 \Gamma((Q\bar{Q} \rightarrow 3gluons)),$ where Q (q) denotes heavy (light) quark, means that each of above listed decays has the branching ratio which is much lower than the branching ratio of the decay into 3 gluons.

QUARKS-2014, June 2-8, 2014, Suzdal – p.32/51



Let us try to understand this suppression in the language of intermediate hadronic states, i.e. in the framework of dispersion approach.

QUARKS-2014, June 2-8, 2014, Suzdal – p.33/51

#### The language of intermediate hadronic states

In this approach a amplitude of a decay under discussion can be represented as the sum over the contributions to the dispersion integral coming from the  $D\bar{D}, D^*\bar{D}+c.c., D^*\bar{D}^*$  etc., intermediate states in the case of the  $J/\psi(1S)$  or  $B\bar{B}, B^*\bar{B}+c.c., B^*\bar{B}^*$  etc., intermediate states in the case of the  $\Upsilon(1S)$ . We do not see a reason for large suppression of each specific contribution. The most probable explanation of the suppression of the decays under consideration is the strong cancellation between the contributions from intermediate states listed above. However, such a cancellation could be broken when a new channel is opening. If so, the energy window may open where imaginary part of the amplitude is appreciable. QUARKS-2014, June 2-8, 2014, Suzdal – p.34/51

 $\psi(3770) \rightarrow PP$ 



We believe that such a situation is realized for the states lying slightly above the production thresholds of open charm and beauty. Hence, the states  $\psi(3770)$  and  $\Upsilon(10580)$  are most promising from the point of view of the idea under consideration. QUARKS-2014, June 2-8, 2014, Suzdal – p.35/51

 $\psi(3770) \rightarrow VP$ 



All told about the  $\psi(3770)$  is transferred to the case of  $\Upsilon(10580) \equiv \Upsilon(4S)$  by means of the replacements  $\psi(3770) \rightarrow$  $\Upsilon(10580), c \rightarrow b, D \rightarrow B, D_s^* \rightarrow B_s^*$  etc.

QUARKS-2014, June 2-8, 2014, Suzdal – p.36/51

 $\psi(3770) 
ightarrow VV$ 





QUARKS-2014, June 2-8, 2014, Suzdal – p.37/51

Mode	$\psi(3770)$	$\Upsilon(10580)$
 $\pi^+\pi^-$	$2\cdot 10^{-6}(7\cdot 10^{-5})$	$8\cdot 10^{-8}(6\cdot 10^{-6})$
$Kar{K}$	$2\cdot 10^{-5}$	$2\cdot 10^{-6}$
$\omega\pi^0$	$2\cdot 10^{-5} (7\cdot 10^{-4})$	$5\cdot 10^{-6}(4\cdot 10^{-4})$
$\omega\eta$	$3\cdot 10^{-4}(1\cdot 10^{-5})$	$3\cdot 10^{-4}(4\cdot 10^{-6})$
$\omega\eta^\prime$	$1\cdot 10^{-4} (7\cdot 10^{-6})$	$2\cdot 10^{-4} (2\cdot 10^{-6})$
$ ho\pi$	$2\cdot 10^{-3}(7\cdot 10^{-5})$	$1\cdot 10^{-3}(2\cdot 10^{-5})$
$ ho\eta$	$1\cdot 10^{-5}(3\cdot 10^{-4})$	$4\cdot 10^{-6}(3\cdot 10^{-4})$
$ ho\eta^\prime$	$7\cdot 10^{-6}(1\cdot 10^{-4})$	$2\cdot 10^{-6}(2\cdot 10^{-4})$
$ ho^+ ho^-$	$3\cdot 10^{-5}(1\cdot 10^{-3})$	$1\cdot 10^{-4}(8\cdot 10^{-3})$
$K^*ar{K}+c.c$	$3\cdot 10^{-4}$	$4\cdot 10^{-4}$
$K^*ar{K}^*$	$7\cdot 10^{-4}$	$3\cdot 10^{-3}$
$J/\psi(1S)+\pi^0$	$8\cdot 10^{-6}(1\cdot 10^{-4})$	-
$J/\psi(1S)+\eta$	$4\cdot 10^{-5}(1\cdot 10^{-6})$	-
$\Upsilon(1S)+\pi^0$	-	$7\cdot 10^{-9} (5\cdot 10^{-7})$
$\Upsilon(1S)+\eta$	-	$2\cdot 10^{-7} (3\cdot 10^{-9})$
3 gluons	$2\cdot 10^{-4}$	$4\cdot 10^{-4}$
total	$4\cdot 10^{-3}(3\cdot 10^{-3})$	$5\cdot 10^{-3}(13\cdot 10^{-3})$

QUARKS-2014, June 2-8, 2014, Suzdal – p.38/51

# Discussion

Calculating the amplitudes, we took into account two suppressing factors: 1) the absorption in final state, the suppressing factor is 1/2; 2) the exponential vertexes of the t-channel exchanges in Figs in the previous slides,  $\exp\{\lambda t\}$  with  $\lambda \approx (1/m_0)^2$ , where  $m_0$  is the mass of the lightest threshold in the t-channel, the effective suppressing factor is 7.

So, if carefully to formulate our results we got a band of predictions for the branching ratios (BR) of decays  $\psi(3770)$  and  $\Upsilon(10580)$ into light (non- $D\bar{D}$  and non- $B\bar{B}$ ) hadrons caused by the intermediate real  $D\bar{D}$  and  $B\bar{B}$ : 1%  $\lesssim$  BR  $\lesssim$  15%. The lower bound is 10 times as large as the branching ratio of annihilation into three gluons.

QUARKS-2014, June 2-8, 2014, Suzdal – p.39/51

How learn the branching ratio  $X(3872) 
ightarrow D^{*0} ar{D}^0 + c.c.$ 

Based on N.N. Achasov and E.V. Rogozina, arxiv:1310.1436 [hep-ph].

The mass spectrum  $\pi^+\pi^- J/\psi(1S)$  in the  $X(3872) \rightarrow \pi^+\pi^- J/\psi(1S)$  decay looks as the ideal Breit-Wigner one, see next Fig.

QUARKS-2014, June 2-8, 2014, Suzdal – p.40/51

### The mass spectrum $\pi^+\pi^-J/\psi(1S)$





a) The Belle data on the invariant  $\pi^+\pi^- J/\psi(1S)$  mass (*m*) distribution. The solid line is our theoretical one with taking into account the Belle energy resolution. b) Our undressed theoretical line.

The mass spectrum  $D^{*0}\overline{D}^0 + c.c.$  in the  $X(3872) \rightarrow D^{*0}\overline{D}^0 + c.c.$  decay looks as the typical resonance threshold enhancement, see next Fig.

QUARKS-2014, June 2-8, 2014, Suzdal – p.41/51

# The mass spectrum $D^{*0}\bar{D}^0+c.c.$



The Belle data on the invariant  $D^{*0}\overline{D}^0 + c.c.$  mass (*m*) distribution. The solid line is our theoretical one with taking into account the Belle energy resolution. a)  $D^{*0} \to D^0 \pi^0$ . b)  $D^{*0} \to D^0 \gamma$ .

If structures in the above channels are manifestation of the same resonance, it is possible to define the branching ratio  $BR(X(3872) \rightarrow D^{*0}\bar{D}^0 + c.c.)$  treating data only about once more decay channel.

# The mass spectrum $D^{*0}\bar{D}^0+c.c.$

We believe that the X(3872) is the axial vector,  $1^{++}$ . In this case the S wave dominates in the  $X(3872) \rightarrow D^{*0}\bar{D}^0 + c.c.$  decay and hence is described by the Lagrangian

$$L(x)=g_AX^\mu\Big(D_\mu(x)ar{D}(x)+ar{D}_\mu(x)D(x)\Big).$$

The width of the  $X 
ightarrow D^{*0} ar{D}^0 + c.c.$  decay

$$\Gamma(X o D^{*0} ar{D}^0 + c.c. \,, \, m) rac{g_A^2}{8\pi} rac{
ho(m)}{m} igg(1 + rac{\mathrm{k}^2}{3m_{D^{*0}}^2}igg) \,,$$

$$ho(m)=rac{2|\mathrm{k}|}{m}=rac{\sqrt{(m^2-m_+^2)(m^2-m_-^2)}}{m^2}, m_{\pm}=m_{D^{*0}}{\pm}m_{D^0}.$$

QUARKS-2014, June 2-8, 2014, Suzdal – p.43/51

# The mass spectrum $D^{*0} ar{D}^0 + c.c.$

$$rac{dBR(X o D^{*0} ar{D}^0 + c.c.\,,\,m)}{dm} = 4 rac{1}{\pi} rac{m^2 \Gamma(X o D^{*0} ar{D}^0,\,m)}{|D_X(m)|^2}$$

The branching ratio of  $X(3872) 
ightarrow D^{*0} ar{D}^0 + c.c.$ 

$$BR(X o D^{*0} ar{D}^0 + c.c.) = 4 rac{1}{\pi} \int_{m_+}^{\infty} rac{m^2 \Gamma(X o D^{*0} ar{D}^0, m)}{|D_X(m)|^2} dm$$

QUARKS-2014, June 2-8, 2014, Suzdal – p.44/51

The mass spectra of X(3872) in non- $D^{*0}ar{D}^0+c.c.$  chanals

In others  $\{i\}$  (non- $D^{*0}\overline{D}^0$ ) channels the X(3872) state is seen as a narrow resonance that is why we write the mass spectrum in the i channel in the form

$$rac{dBR(X 
ightarrow i\,,\,m)}{dm} = 2 rac{1}{\pi} rac{m_X^2 \, \Gamma_i}{|D_X(m)|^2}\,,$$

where  $\Gamma_i$  is the width of the X(3872) 
ightarrow i decay.

The branching ratio of X(3872) 
ightarrow i

$$BR(X
ightarrow i)=2rac{1}{\pi}\int_{m_0}^\infty rac{m_X^2\Gamma_i}{|D_X(m)|^2}dm\,,$$

where  $m_0$  is the threshold of the i state.

QUARKS-2014, June 2-8, 2014, Suzdal – p.45/51

# The inverse propagator $D_X(m)$

 $D_X(m) = m_X^2 - m^2 + Re(\Pi_X(m_X)) - \Pi_X(m) - \imath m_X \Gamma$ , where  $\Gamma = \Sigma \Gamma_i$  is the total width of the X(3872) decay into all non- $D^{*0}\bar{D}^0$  channels.

When 
$$m_+ \leq m$$
,  
 $\Pi_X(m) = rac{g_A^2}{8\pi^2} \left\{ rac{(m^2 - m_+^2)}{m^2} rac{m_-}{m_+} \ln rac{m_{D^{*0}}}{m_{D^0}} + 
ho(m) \left[ \imath \pi + \ln rac{\sqrt{m^2 - m_-^2} - \sqrt{m^2 - m_+^2}}{\sqrt{m^2 - m_-^2} + \sqrt{m^2 - m_+^2}} \right] 
ight\}.$ 

In other areas of m ( $m_{-} \leq m \leq m_{+}$ ,  $m \leq m_{-}$  and  $m^{2} \leq 0$ ), the function  $\Pi_{X}(m)$  is defined by analytic continuation.

QUARKS-2014, June 2-8, 2014, Suzdal – p.46/51

# **Unitarity, Fitting Data**

Our branching ratios satisfy unitarity

$$1 = BR(X \rightarrow D^{*0}\overline{D}^0 + c.c.) + \sum_i BR(X \rightarrow i).$$

Fitting the Belle data, we take into account the Belle results that  $m_X = 3871.84 \text{ MeV} = m_{D^{*0}} + m_{D^0} = m_+$  and  $\Gamma_{X(3872)} < 1.2 \text{ MeV}$  90%CL that corresponds to  $\Gamma < 1.2 \text{ MeV}$ , which controls the width of the X(2872) signal in the  $\pi^+\pi^-J/\psi(1S)$  channel and in every non- $D^{*0}\bar{D}^0$  channel.

The results of our fit are in the Table.

QUARKS-2014, June 2-8, 2014, Suzdal – p.47/51

### **Results and Discusion**

TABLE. Results of the analysis of the Belle data .

$$\begin{split} B_{seen} &= BR(X \to D^{*0}\bar{D}^0 + c.c.; m \leq 3891.84 \, \text{MeV}), \\ B &= BR(X \to D^{*0}\bar{D}^0 + c.c.), \\ B(r)_{seen} &= \sum_i BR(X \to i; \, 3851.84 \leq m \leq 3891.84 \, \text{MeV}), \\ \text{Sum} &= BR(X \to D^{*0}\bar{D}^0 + c.c.) + \sum_i BR(X \to i) = 1. \end{split}$$

$\overline{\Gamma}$	$g_A^2/8\pi$	$\chi^2/Ndf$	$\mathcal{B}_{seen}$	B	$\mathcal{B}(r)_{seen}$	Sum
$1.2_{-0.5}$	$0.9^{+4}_{-0.5}$	44/42	$0.5\substack{+0.1\-0.3}$	$0.8\substack{+0.2\-0.2}$	$0.2\substack{+0.2 \\ -0.2}$	1

The current statistics is not sufficient for serious conclusions.Nevertheless, one can state that our results are consist with experiment.

QUARKS-2014, June 2-8, 2014, Suzdal – p.48/51

### **Results and Discusion**

#### Really, in view of

 $\begin{array}{l} BR(B\to X(3872)K)\times BR(X(3872)\to D^{*0}\bar{D}^0)=\\ (0.80\pm 0.20\pm 0.1)\times 10^{-4}, \end{array}$ 

 $\begin{array}{l} BR(B^+ \to X(3872)K^+) \times BR(X(3872) \to \pi^+\pi^-J/\psi(1S)) = \\ (8.61 \pm 0.82 \pm 0.52) \times 10^{-6}, \end{array}$ 

$$\begin{split} BR(B^+ \to X(3872)K^+) \times BR(X(3872) \to \pi^+\pi^-\pi^0 J/\psi(1S)) = \\ (0.6 \pm 0.2 \pm 0.1) \times 10^{-5}, \end{split}$$

 $\begin{array}{l} BR(B^+ \to X(3872)K^+) \times BR(X(3872) \to \gamma J/\psi(1S)) = \\ (1.78^{+0.48}_{-0.44} \pm 0.12) \times 10^{-6} \end{array}$ 

it follows that  $BR(X \to D^{*0}\bar{D}^0 + c.c.; m \le 3892 \,\text{MeV})$  is a few times as large as the sum of all non- $D^{*0}\bar{D}^0$  known branching ratios.

QUARKS-2014, June 2-8, 2014, Suzdal – p.49/51

# Discusion

So, when fitting the  $X(3872) \rightarrow D^{*0}\bar{D}^0$  data and data for any X(3872) decay into non- $D^{*0}\bar{D}^0$  state,  $X(3872) \rightarrow i$ , we find  $\Gamma$  and  $g_A^2/4\pi$ , which define  $BR(X(3872) \rightarrow D^{*0}\bar{D}^0 + c.c.)$ . Generally speaking, we don't need to know  $BR(X(3872) \rightarrow i)$ .

Our approach can serve as the guide in selection of theoretical models for the X(3872) resonance. Indeed,

 $3871.68~{
m MeV} < M_X < 3871.95~{
m MeV}$  and  $\Gamma_{M_X} = \Gamma < 1.2~{
m MeV}.$ 

Let  $g_A^2/4\pi < 0.2 \text{ GeV}^2$  that does not contradict current experiment, generally speaking. But then

 $BR(X 
ightarrow D^{*0} ar{D}^0 + c.c.\,; m \leq 3891.84\, {
m MeV}) = BR_{seen} < 0.3$  ,

that is, unknown decays of X(3872) into non- $D^{*0}\bar{D}^0$  states are considerable or dominant.

QUARKS-2014, June 2-8, 2014, Suzdal – p.50/51



This work was supported in part by RFBR, Grant No. 13-02-00039, and Interdisciplinary project No. 102 of Siberian division of RAS.

THANK YOU

QUARKS-2014, June 2-8, 2014, Suzdal – p.51/51